

GENERALIZED VOLUME CONJECTURE AND THE A-POLYNOMIALS: THE NEUMANN–ZAGIER POTENTIAL FUNCTION AS A CLASSICAL LIMIT OF QUANTUM INVARIANT

KAZUHIRO HIKAMI

Abstract. We study quantum invariant $Z_\gamma(\mathcal{M})$ for cusped hyperbolic 3-manifold $\mathcal{M}$. We construct this invariant based on oriented ideal triangulation of $\mathcal{M}$ by assigning to each tetrahedron the quantum dilogarithm function, which is introduced by Faddeev in studies of the modular double of the quantum group. Following Thurston and Neumann–Zagier, we deform a complete hyperbolic structure of $\mathcal{M}$, and correspondingly we define quantum invariant $Z_\gamma(\mathcal{M}_u)$. This quantum invariant is shown to give the Neumann–Zagier potential function in the classical limit $\gamma \rightarrow 0$, and the A-polynomial can be derived from the potential function. We explain our construction by taking examples of 3-manifolds such as complements of hyperbolic knots and punctured torus bundle over the circle.

1. I

Since the quantum invariant of knots/links and 3-manifolds as a generalization of the Jones polynomial [25] is constructed by Witten [53] by use of the Chern–Simons path integral, studies on quantum invariants have been much developed. Recently geometrical interpretations of the quantum invariants have received interests since Kashaev observed an intriguing relationship [29] between the hyperbolic volume and his knot invariant, which is later identified with a specific value of the N -colored Jones polynomial $J_{\mathcal{K}}(N; e^{2\pi i/N})$ [41] (here the N -colored Jones polynomial is normalized to be $J_{\text{unknot}}(N; q) = 1$). Namely the hyperbolic volume of the knot complement $S^3 \setminus \mathcal{K}$ is conjectured to dominate the asymptotics of the invariant $J_{\mathcal{K}}(N; e^{2\pi i/N})$ in the large- N limit $N \rightarrow \infty$,

$$\lim_{N \rightarrow \infty} \frac{2\pi}{N} \log |J_{\mathcal{K}}(N; e^{2\pi i/N})| = \text{Vol}(S^3 \setminus \mathcal{K}) \quad (1.1)$$

This “volume conjecture” is generalized for other values (near the N -th root of unity) of the N -colored Jones polynomial [18] (see also Refs. 39, 40), and a relationship with the A-polynomial is conjectured; when we define b by

$$b = -\frac{d}{da} \lim_{\substack{N, k \rightarrow \infty \\ N/k = a}} \frac{1}{k} \log J_{\mathcal{K}}(N; e^{2\pi i/k}) \quad (1.2)$$

Date: April 4, 2006.

the pair $(e^b, -e^{ia})$ is a zero locus of the A-polynomial for knot $\mathcal{K}$. This is checked numerically for twist knots [22].

The A-polynomial is defined as an algebraic curve of eigenvalues of the $SL(2; \mathbb{C})$ representation of the boundary torus of knot $\mathcal{K}$ [9] (see also Ref. 10). This can be computed from triangulation of the knot complement $\mathcal{M} = S^3 \setminus \mathcal{K}$ into ideal tetrahedra in the hyperbolic space $\mathbb{H}^3$ once the fundamental group has a irreducible representation ρ into $PSL(2; \mathbb{C})$ which is identified with the orientation-preserving isometries of $\mathbb{H}^3$;

$$\rho : \pi_1(\mathcal{M}) \rightarrow PSL(2; \mathbb{C}) \simeq \text{Isom}^+(\mathbb{H}^3)$$

Up to conjugation, the meridian μ and the longitude of the boundary torus of $\mathcal{K}$ have

$$\rho(\mu) = \begin{pmatrix} m & * \\ 0 & 1/m \end{pmatrix}$$

$$\rho(\lambda) = \begin{pmatrix} \ell & * \\ 0 & 1/\ell \end{pmatrix}$$

Another geometrical aspect of the N -colored Jones polynomial $J_{\mathcal{K}}(N; q)$ as a relationship with the A-polynomial is proposed as “AJ conjecture” [17]; recursion relation of the colored Jones polynomial with respect to N is conjectured to be related to the A-polynomial $A_{\mathcal{K}}(\ell, m)$. This conjecture is proved for the torus knots [23] and the 2-bridge knots [35].

In this paper, we introduce quantum invariant for cusped hyperbolic manifold $\mathcal{M}$ and its deformation $\mathcal{M}_u$ à la Thurston [50] following Refs. 19–21, and study a classical limit thereof. Based on a triangulation of cusped 3-manifold $\mathcal{M}$, we define quantum invariant $Z_{\gamma}(\mathcal{M}_u)$ by assigning Faddeev’s quantum dilogarithm function to each oriented ideal tetrahedron. Originally Kashaev introduced his invariant $J_{\mathcal{K}}(N; e^{2\pi i/N})$ for triangulated 3-manifolds, although the R -matrix construction is developed subsequently [28]. He studied Faddeev’s quantum dilogarithm function when q is a root of unity [13, 26] (see also Ref. 2), and assigning the quantum dilogarithm function to ideal tetrahedron he defined invariant [27, 28]. In this sense, our invariant $Z_{\gamma}(\mathcal{M})$ for $\mathcal{M}$ with complete hyperbolic structure can be regarded as a non-compact $U_q(sl(2; \mathbb{R}))$ analogue of the Kashaev invariant, and an asymptotic behavior of $Z_{\gamma}(S^3 \setminus \mathcal{K})$ in the limit $\gamma \rightarrow 0$ should coincide with that of the Kashaev invariant $J_{\mathcal{K}}(N; e^{2\pi i/N})$ in the limit $N \rightarrow \infty$.

This paper is organized as follows. In Section 2 we recall definitions of quantum dilogarithm function, and we discuss properties of this function. In Section 3, we shall reveal that the three-dimensional hyperbolic geometry naturally arises from the quantum dilogarithm function in the classical limit $\gamma \rightarrow 0$ as was clarified in Ref. 19. Then we define the quantum invariant $Z_{\gamma}(\mathcal{M}_u)$ for a deformation of complete hyperbolic cusped 3-manifold $\mathcal{M}$. Based on a triangulation of

$\mathcal{M}$, we construct the invariant by assigning quantum dilogarithm function to oriented ideal tetrahedron. We discuss that the Neumann–Zagier potential function appears in a classical limit of $Z_\gamma(\mathcal{M})$. In Section 4, we take several examples of cusped hyperbolic manifolds such as complements of hyperbolic knots and punctured torus bundle over the circle, and explain our assertion in detail. We shall also give a list for other manifolds in Appendix. The last section is devoted to conclusions and discussions.

2. Q D F

We define a function $\Phi_\gamma(\varphi)$ by an integral form following Ref. 12. We set $\gamma \in \mathbb{R}$, and for $|\operatorname{Im} \varphi| < \pi$, we define

$$\Phi_\gamma(\varphi) = \exp \left(\int_{\mathbb{R}+i0} \frac{e^{-i\varphi x}}{4 \sinh(\gamma x) \sinh(\pi x)} \frac{dx}{x} \right) \quad (2.1)$$

The Faddeev integral (2.1), which we call the quantum dilogarithm function, is also related to the double sine function [31, 34, 48], the hyperbolic gamma function [1, 46] and the quantum exponential function [54]. We see that the integral $\Phi_\gamma(\varphi)$ has a duality,

$$\Phi_{\frac{\pi^2}{\gamma}}(\varphi) = \Phi_\gamma\left(\frac{\gamma}{\pi} \varphi\right) \quad (2.2)$$

and that it satisfies the inversion relation,

$$\Phi_\gamma(\varphi) \cdot \Phi_\gamma(-\varphi) = \exp \left(-\frac{1}{2i\gamma} \left(\frac{\varphi^2}{2} + \frac{\pi^2 + \gamma^2}{6} \right) \right) \quad (2.3)$$

The Faddeev integral satisfies the difference equations;

$$\frac{\Phi_\gamma(\varphi + i\gamma)}{\Phi_\gamma(\varphi - i\gamma)} = \frac{1}{1 + e^\varphi} \quad (2.4a)$$

$$\frac{\Phi_\gamma(\varphi + i\pi)}{\Phi_\gamma(\varphi - i\pi)} = \frac{1}{1 + e^{\frac{\pi}{\gamma}\varphi}} \quad (2.4b)$$

Due to these relations, the integral $\Phi_\gamma(\varphi)$ defined in (2.1) is analytically continued to $\varphi \in \mathbb{C}$, and we see that

$$\text{zeros of } (\Phi_\gamma(\varphi))^{\pm 1} = \left\{ \varphi = \mp i((2m+1)\gamma + (2n+1)\pi) \mid m, n \in \mathbb{Z}_{\geq 0} \right\} \quad (2.5)$$

The most important properties of the Faddeev integral is that it fulfills the pentagon identity [12, 13]

$$\Phi_\gamma(\hat{p}) \Phi_\gamma(\hat{q}) = \Phi_\gamma(\hat{q}) \Phi_\gamma(\hat{p} + \hat{q}) \Phi_\gamma(\hat{p}) \quad (2.6)$$

where $\hat{p}$ and $\hat{q}$ are the canonically conjugate operators satisfying the Heisenberg commutation relation,

$$[\hat{p}, \hat{q}] = \hat{p}\hat{q} - \hat{q}\hat{p} = -2i\gamma \quad (2.7)$$

By this commuting relation, we call a limit $\gamma \rightarrow 0$ a classical limit.

We use $\mathbf{V}$ as the momentum space $|p\rangle$ with $p \in \mathbb{R}$ which is an eigenstate of the momentum operator;

$$\hat{p}|p\rangle = p|p\rangle. \quad (2.8)$$

A reason of the *quantum* dilogarithm function reveals when we take a *classical* limit $\gamma \rightarrow 0$. In this limit, the Faddeev integral reduces to

$$\Phi_\gamma(\varphi) \sim \exp\left(\frac{1}{2i\gamma} \text{Li}_2(-e^\varphi)\right) \quad (2.9)$$

where $\text{Li}_2(x)$ denotes the Euler dilogarithm function defined by (see, *e.g.*, Refs. 32, 36)

$$\text{Li}_2(x) = \sum_{n=1}^{\infty} \frac{x^n}{n^2} \quad (2.10)$$

where $|x| \leq 1$. For $x \in \mathbb{C}$, we use the integral form

$$\text{Li}_2(x) = - \int_0^x \log(1-s) \frac{ds}{s}$$

where the branch of $\log(1-s)$ is on $\mathbb{C} \setminus [1, \infty)$ for which $\log(1-0) = 0$. See that the inversion relation (2.3) of the quantum dilogarithm function $\Phi_\gamma(\varphi)$ gives that of the Euler dilogarithm function as

$$\text{Li}_2(-e^x) + \text{Li}_2(-e^{-x}) + \frac{x^2}{2} + \frac{\pi^2}{6} = 0$$

We note that the Fourier transformation of the Faddeev integral can be computed as follows [5, 14, 31, 54];

$$\frac{1}{\sqrt{4\pi\gamma}} \int_{\mathbb{R}} dy \Phi_\gamma(y) e^{\frac{1}{2i\gamma}xy} = \Phi_\gamma(-x + i\pi + i\gamma) e^{\frac{1}{2i\gamma}\left(\frac{x^2}{2} - \frac{1}{2}\pi\gamma - \frac{\pi^2 + \gamma^2}{6}\right)} \quad (2.11)$$

$$\frac{1}{\sqrt{4\pi\gamma}} \int_{\mathbb{R}} dy \frac{1}{\Phi_\gamma(y)} e^{\frac{1}{2i\gamma}xy} = \frac{1}{\Phi_\gamma(x - i\pi - i\gamma)} e^{-\frac{1}{2i\gamma}\left(\frac{x^2}{2} - \frac{1}{2}\pi\gamma - \frac{\pi^2 + \gamma^2}{6}\right)} \quad (2.12)$$

To see a relationship between the integral $\Phi_\gamma(\varphi)$ with geometry, we define the S -operator acting on $\mathbf{V} \otimes \mathbf{V}$ by

$$S_{1,2} = e^{\frac{1}{2i\gamma} \hat{q}_1 \hat{p}_2} \Phi_\gamma(\hat{p}_1 + \hat{q}_2 - \hat{p}_2) \quad (2.13)$$

Here the Heisenberg operators $\hat{p}_j$ and $\hat{q}_j$ act on the j -th vector space of $\mathbf{V} \otimes \mathbf{V}$, i.e. $\hat{p}_1 = \hat{p} \otimes 1$, $\hat{p}_2 = 1 \otimes \hat{p}$, and so on. Then the pentagon identity (2.6) can be rewritten in a compact form;

$$S_{2,3} S_{1,2} = S_{1,2} S_{1,3} S_{2,3} \quad (2.14)$$

where $S_{j,k}$ acts on the j - and k -th spaces of $\mathbf{V} \otimes \mathbf{V} \otimes \mathbf{V}$. Matrix elements can be computed by use of (2.11) and (2.12);

$$\begin{aligned} \langle p_1, p_2 | S_{1,2} | p'_1, p'_2 \rangle \\ = \frac{1}{\sqrt{4\pi\gamma}} \delta(p_1 + p_2 - p'_1) \cdot \Phi_\gamma(p'_2 - p_2 + i\pi + i\gamma) e^{\frac{1}{2i\gamma} \left(-\frac{\pi^2 + \gamma^2}{6} - \frac{\gamma\pi}{2} + p_1(p'_2 - p_2) \right)} \end{aligned} \quad (2.15a)$$

$$\begin{aligned} \langle p_1, p_2 | S_{1,2}^{-1} | p'_1, p'_2 \rangle \\ = \frac{1}{\sqrt{4\pi\gamma}} \delta(p_1 - p'_1 - p'_2) \frac{1}{\Phi_\gamma(p_2 - p'_2 - i\pi - i\gamma)} e^{\frac{1}{2i\gamma} \left(\frac{\pi^2 + \gamma^2}{6} + \frac{\gamma\pi}{2} - p'_1(p_2 - p'_2) \right)} \end{aligned} \quad (2.15b)$$

In the classical limit $\gamma \rightarrow 0$, we find by use of (2.9) that the S -operators (2.15) reduce to

$$\langle p_1, p_2 | S_{1,2} | p'_1, p'_2 \rangle \sim \delta(p_1 + p_2 - p'_1) \cdot \exp \left(-\frac{1}{2i\gamma} V(p'_2 - p_2, p_1) \right) \quad (2.16a)$$

$$\langle p_1, p_2 | S_{1,2}^{-1} | p'_1, p'_2 \rangle \sim \delta(p_1 - p'_1 - p'_2) \cdot \exp \left(\frac{1}{2i\gamma} V(p_2 - p'_2, p'_1) \right) \quad (2.16b)$$

where we have defined the function $V(x, y)$ by

$$V(x, y) = \frac{\pi^2}{6} - \text{Li}_2(e^x) - xy. \quad (2.17)$$

We see that the function $V(x, y)$ satisfies partial differential equations;


$$V(x, y) = L(1 - e^x) + \frac{1}{2} \left(x \frac{\partial V(x, y)}{\partial x} + y \frac{\partial V(x, y)}{\partial y} \right) \quad (2.18)$$

$$\text{Im } V(x, y) = D(1 - e^x) + \log |e^x| \cdot \text{Im} \left(\frac{\partial}{\partial x} V(x, y) \right) + \log |e^y| \cdot \text{Im} \left(\frac{\partial}{\partial y} V(x, y) \right) \quad (2.19)$$

Here the Rogers dilogarithm $L(z)$ and the Bloch–Wigner function $D(z)$ are respectively defined in terms of the Euler dilogarithm function (2.10) by

$$L(z) = \text{Li}_2(z) + \frac{1}{2} \log z \log(1 - z) \quad (2.20)$$

$$D(z) = \text{Im } \text{Li}_2(z) + \arg(1 - z) \cdot \log |z| \quad (2.21)$$



both of which fulfill the pentagon identity (see, *e.g.*, Ref. 36);

$$L(z) - L(w) + L\left(\frac{w}{z}\right) - L\left(\frac{1-z^{-1}}{1-w^{-1}}\right) + L\left(\frac{1-z}{1-w}\right) = \frac{\pi^2}{6} \quad (2.22)$$

$$D(z) - D(w) + D\left(\frac{w}{z}\right) - D\left(\frac{1-z^{-1}}{1-w^{-1}}\right) + D\left(\frac{1-z}{1-w}\right) = 0 \quad (2.23)$$


3. Q I P F

The result that the S -operator satisfies the pentagon identity (2.14) indicates that we can assign (oriented) tetrahedron to each S -operator. See Refs. 7, 30 (also Ref. 8) for another interpretation as a quantization of the Teichmüller theory. In the classical limit $\gamma \rightarrow 0$, imaginary part of the S -operator gives the Bloch–Wigner function $D(z)$ which denotes the hyperbolic volume of the ideal tetrahedron $\Delta(z)$ with totally geodesic faces which has a modulus z (see, *e.g.*, Refs. 3, 38, 50). Extra terms in (2.19) including partial differentials of $V(x, y)$ can be neglected when we assume a saddle point condition as will be discussed later, and we can assign the oriented hyperbolic ideal tetrahedra to these S -operators as follows [19];

$$\langle a, b | S | c, d \rangle = \text{Diagram} = \quad (3.1)$$

$$\langle a, b | S^{-1} | c, d \rangle = \text{Diagram} = \quad (3.2)$$

Namely in the momentum space representation we assign four momentum parameters $\{a, b, c, d\}$ to faces of tetrahedron. This differs from the (quantum) $6j$ symbol as a solution of the pentagon identity [33, 45], which was used to define the topological (quantum) gravity.



F 1. Triangle with vertices 0, 1, and z in $\mathbb{C}$. Here we set $z[1] = z$, $z[2] = 1 - \frac{1}{z}$, and $z[3] = \frac{1}{1-z}$.

Here both ideal tetrahedra have the modulus $z = e^{d-b}$, and we mean that the dihedral angles are given by

$$z[1] = z = e^{d-b}$$

$$z[2] = 1 - \frac{1}{z}$$

$$z[3] = \frac{1}{1-z}$$

The cross sections by the horosphere is similar to the triangle in $\mathbb{C}$ with vertices 0, 1, and z (see Fig. 1), and we have $z[1]z[2]z[3] = -1$. See that the opposite edges of tetrahedra have the same dihedral angles. Then the hyperbolic volume of the ideal tetrahedron $\Delta(z)$ with modulus z is given by [50]


$$\begin{aligned} \text{Vol}(\Delta(z)) &= D(z) \\ &= \mathcal{L}(\arg(z[1])) + \mathcal{L}(\arg(z[2])) + \mathcal{L}(\arg(z[3])) \end{aligned} \quad (3.3)$$

where $\mathcal{L}(\theta)$ is the Lobachevsky function defined by

$$\mathcal{L}(\theta) = \frac{1}{2} \sum_{n=1}^{\infty} \frac{\sin(2n\theta)}{n^2} \quad (3.4)$$

Any hyperbolic cusped 3-manifold $\mathcal{M}$ (for simplicity we assume that the manifold $\mathcal{M}$ has one cusp in this paper) can be ideally triangulated, and it is constructed from finite number of the oriented ideal tetrahedra in (3.1) and (3.2). Other types of oriented ideal tetrahedra, such as one that has a face with loop, are prohibited. So our triangulation sometimes differ from the canonical triangulation used in computer programs, such as SnapPea [52], Knotscape [24], and Snap [11].

To construct 3-manifolds from these ideal tetrahedra, we glue each face to match orientations of edges. Every faces have no loops, so there is no ambiguity in gluing faces together once two faces to be glued are fixed. Furthermore, one finds that in-state $|p\rangle$ can glue only with



F 2. Pentagon identity (2.14) is interpreted as the Pachner move.

out-states $\langle p|$, and that we cannot glue two in-states or out-states together. As suggested by the momentum representation of the pentagon identity, gluing two faces, which have same label p , can be formulated by an integration with respect to p , $\int_{\mathbb{R}} dp |p\rangle \langle p|$.

By these identifications of the S -operators as the oriented ideal tetrahedra, the pentagon identity (2.14) is identified with the Pachner move as is depicted in Fig. 2; oriented polytope with 5 vertices can be decomposed into 2 tetrahedra with a face in common, or into 3 tetrahedra with an edge in common. In case of decomposing into 3 tetrahedra, we need a hyperbolic consistency condition around the common edge. This condition coincides with a saddle point condition of the right hand side of (2.14) after substituting (2.15) and taking a classical limit $\gamma \rightarrow 0$. One can see a coincidence for any other type of gluings such as other choices of orientations in the Pachner move between the hyperbolic consistency condition and the saddle point equations [19].

Once triangulation is given and we know how to glue faces together, we can naturally define the quantum invariant for hyperbolic cusped 3-manifold $\mathcal{M}$ based on the S -operator by

$$Z_\gamma(\mathcal{M}) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}) \delta_G(\mathbf{p}) \prod_{i=1}^M \langle p_{2i-1}^{(-)}, p_{2i}^{(-)} | S^{\varepsilon_i} | p_{2i-1}^{(+)}, p_{2i}^{(+)} \rangle \quad (3.5)$$

where $\mathbf{p}$ denotes a set of variables $(p_1^{(\pm)}, p_2^{(\pm)}, \dots, p_{2M}^{(\pm)})$, and $\varepsilon_i = \pm 1$ depending on an orientation of tetrahedron. We set M as the number of ideal tetrahedra. The condition $\delta_G(\mathbf{p})$ determines how to glue faces together (“G” stands for “gluing”). Every faces with same momentum mean to

be glued together, and the fact that in-states can be glued only with out-states indicates that the gluing condition $\delta_G(\mathbf{p})$ is a product of $\delta(p_j^{(-)} - p_k^{(+)})$

We need another geometrical condition $\delta_C(\mathbf{p})$ to define invariant besides a way how to glue faces (“C” stands for “completeness”) [21]. We can draw a developing map from an ideal triangulation of 3-manifold, and we need to read off a hyperbolic complete condition. This condition can be written as a constraint for $\mathbf{p}$ by identifications (3.1) and (3.2). By construction, the partition function $Z_\gamma(\mathcal{M})$ is invariant under the Pachner move (Fig. 2), and it is essentially the invariant constructed in Ref. 19.

As was studied by Thurston [51], *deformation* of the hyperbolic structure on the manifold $\mathcal{M}$ can be holomorphically parametrized by a parameter u in a neighborhood of completeness condition $u = 0$. The parameter u is the logarithm of the eigenvalue of the meridian by the holonomy representation, and we set

$$m = e^u \quad (3.6)$$

Correspondingly we call such manifold $\mathcal{M}_u$ which is no more complete, and define the invariant by

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \delta_G(\mathbf{p}) \prod_{i=1}^M \langle p_{2i-1}^{(-)}, p_{2i}^{(-)} | S^{\varepsilon_i} | p_{2i-1}^{(+)}, p_{2i}^{(+)} \rangle \quad (3.7)$$

Here the condition $\delta_C(\mathbf{p}; u)$ follows from that the meridian has a holonomy (3.6). The invariant (3.5) for $\mathcal{M}$ with complete hyperbolic structure follows from (3.7) by assuming a completeness $u = 0$ of the hyperbolic structure of $\mathcal{M}_u$;

$$Z_\gamma(\mathcal{M}) = Z_\gamma(\mathcal{M}_{u=0}) \quad (3.8)$$

because

$$\delta_C(\mathbf{p}) = \delta_C(\mathbf{p}; u = 0) \quad (3.9)$$

We now study a classical limit of our quantum invariant $Z_\gamma(\mathcal{M}_u)$. When we take a limit $\gamma \rightarrow 0$ by use of (2.16), the quantum invariant defined by (3.7) becomes

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) &\sim \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \delta_G(\mathbf{p}) \left[\prod_{i=1}^M \delta(p_{2i-1}^{(\varepsilon_i)} - p_{2i-1}^{(-\varepsilon_i)} - p_{2i}^{(-\varepsilon_i)}) \right] \\ &\quad \times \exp\left(\frac{i}{2\gamma} \sum_{i=1}^M \varepsilon_i V(p_{2i}^{(\varepsilon_i)} - p_{2i}^{(-\varepsilon_i)}, p_{2i-1}^{(-\varepsilon_i)})\right) \\ &= \int_{\mathbb{R}} d\mathbf{x} \exp\left(\frac{1}{2i\gamma} \Phi_{\mathcal{M}}(\mathbf{x}; u)\right) \end{aligned} \quad (3.10)$$

Here in the last equality for our convention we have re-parametrized variables $\mathbf{p}$ with $\mathbf{x} = (x_1, x_2, \dots, x_{M-1})$ after incorporating constraints written in terms of delta functions.

The integral (3.10) could be evaluated by the saddle point method as we have taken a classical limit $\gamma \rightarrow 0$. The saddle point condition for variables $\mathbf{x}$ is

$$\frac{\partial}{\partial x_i} \Phi_{\mathcal{M}}(\mathbf{x}; u) = 0 \quad (3.11)$$

As was extensively studied in Ref. 19, these conditions coincide with hyperbolic consistency conditions around edges when we glue oriented tetrahedra together. By construction the variable u denotes the meridian of cusp in this classical limit, and the complete hyperbolic structure is realized by setting $u = 0$. To conclude, the function $\Phi_{\mathcal{M}}(\mathbf{x}; u)$ defined by a classical limit (3.10) of the quantum invariant $Z_{\gamma}(\mathcal{M}_u)$ is nothing but the Neumann–Zagier potential function [44, 55].

As a result, differential of the potential function with respect to the deformation parameter u gives

$$\frac{\partial}{\partial u} \Phi_{\mathcal{M}}(\mathbf{x}; u) = 2v \quad (3.12)$$

where v is related to the eigenvalue of the longitude by the holonomy representation

$$\ell = e^v \quad (3.13)$$

Variables x_i can be solved from the hyperbolic consistency equation (3.11) as a function of u , and we can regard the potential function $\Phi_{\mathcal{M}}(\mathbf{x}; u)$ as a function of u ; $\Phi_{\mathcal{M}}(u) = \Phi_{\mathcal{M}}(\mathbf{x}; u)|_{(3.11)}$. Then we can rewrite (3.12) by

$$\frac{d}{du} \lim_{\gamma \rightarrow 0} i \gamma \log Z_{\gamma}(\mathcal{M}_u) = v \quad (3.14)$$

We note that our variables (u, v) differs from those in Ref. 44; when we denote their variables $(u_{\text{NZ}}, v_{\text{NZ}})$, we have

$$(u, v) = \left(\frac{u_{\text{NZ}}}{2}, \frac{v_{\text{NZ}}}{2} + \pi i \right) \quad (3.15)$$

Hereafter we also use the function $V_{\mathcal{M}}(\mathbf{x}; m)$ defined by

$$V_{\mathcal{M}}(x_1, x_2, \dots; m) = \Phi_{\mathcal{M}}(\log x_1, \log x_2, \dots; u = \log m) \quad (3.16)$$

As seen from (2.18) and (2.19), the potential function $\Phi_{\mathcal{M}}(\mathbf{x}; u)$ under saddle point conditions (3.11) becomes a sum of the Rogers dilogarithm functions. We recall here the Bloch invariant studied in Refs. 42, 43. The Bloch invariant $\beta(\mathcal{M})$ is defined for finite volume hyperbolic 3-manifold $\mathcal{M}$ as

$$\beta(\mathcal{M}) = \sum_{i=1}^M [z_i] \quad (3.17)$$

where $[z]$ satisfies the Bloch group

$$[z] - [w] + \left[\frac{w}{z} \right] - \left[\frac{1 - z^{-1}}{1 - w^{-1}} \right] + \left[\frac{1 - z}{1 - w} \right] = 0 \quad (3.18)$$

The Bloch regulator map ρ gives [43]

$$\rho(\beta(\mathcal{M})) = \text{Vol}(\mathcal{M}) + i \text{CS}(\mathcal{M}) \quad (3.19)$$

where CS denotes the Chern–Simons invariant (see Ref. 37 for definition of the Chern–Simons invariant for a case of cusped manifolds). As identities (2.18)–(2.19) show that the S -operator reduces to the Rogers dilogarithm function (or the Bloch–Wigner function) in the saddle point, and that they satisfy the pentagon identities (2.22) and (2.23), we can interpret our invariant $Z_\gamma(\mathcal{M})$ as a quantization of the Bloch invariant.

Generally we have many saddle points as algebraic solutions of a set of equations (3.11). Among them, a solution which has the largest absolute value dominates an asymptotics of the quantum invariant $Z_\gamma(\mathcal{M})$. Combining this with a fact that our quantum invariant $Z_\gamma(\mathcal{M})$ may be regarded as a quantization of the Bloch invariant, we should have

$$\lim_{\gamma \rightarrow 0} 2\gamma \log(Z_\gamma(\mathcal{M})) = \text{Vol}(\mathcal{M}) + i \text{CS}(\mathcal{M}) \quad (3.20)$$

as a variant of the volume conjecture [19]. Although, there still remains an ambiguity of branch in complex plane in an actual computation.

The Neumann–Zagier potential function $\Phi_{\mathcal{M}}(\mathbf{x}; u)$ has much information on geometry of manifold. One of the properties is a relationship with the A-polynomial defined in Ref. 9. When the manifold $\mathcal{M}$ is a complement of knot $S^3 \setminus \mathcal{K}$, the A-polynomial $A_{\mathcal{K}}(\ell, m)$ of $\mathcal{K}$ as an algebraic equation of ℓ and m can be given by using the Gröbner base or resultant theory to eliminate x_i from a set of equations, (3.11) and (3.12). So the pair $(\ell, m) = (e^v, e^u)$ defined from (3.14) is a zero locus of the A-polynomial $A_{\mathcal{K}}(\ell, m)$. This result should be comparable with a conjecture (1.2), and the invariant $Z_\gamma(\mathcal{M}_u)$ is a *non-compact* generalization of the Jones–Witten invariant.

The A-polynomial has following properties [9, 10];

- Polynomial $A_{\mathcal{K}}(\ell, m)$ is an integer polynomial, and it contains only even powers of m .
- Up to powers of ℓ and m , we have

$$A_{\mathcal{K}}(\ell, m) = A_{\mathcal{K}}(1/\ell, 1/m) \quad (3.21)$$

- If $\mathcal{K}$ and $\mathcal{K}'$ are mirror images, then

$$A_{\mathcal{K}}(\ell, m) = A_{\mathcal{K}'}(1/\ell, m) \quad (3.22)$$

- With additional property that every closed incompressible surface embedded in $S^3 \setminus \mathcal{K}$ is parallel to the boundary torus, we have

$$A_{\mathcal{K}}(\ell, \pm 1) = n (\ell + 1)^{\alpha_+} (\ell - 1)^{\alpha_-} \ell^{\beta} \quad (3.23)$$

with non-zero integer n .

- Slopes of edges of the Newton polygon of $A_{\mathcal{K}}(\ell, m)$ are boundary slopes of $\mathcal{K}$.
- Coefficients of terms in the corners of the Newton polygons of $A_{\mathcal{K}}(\ell, m)$ are ± 1 .

Another property of the Neumann–Zagier potential function $\Phi_{\mathcal{M}}(u)$ associated to cusped manifold $\mathcal{M}$ is the volume of the Dehn surgered manifold. The (p, q) -hyperbolic Dehn surgery of $\mathcal{M}$, where (p, q) is a pair of coprime integers, is performed by gluing back a solid torus with cusp of $\mathcal{M}$, where the surgery data satisfy [50]

$$p u + q v = \pi i \quad (3.24)$$

Then for the core c of solid torus, we have

$$\text{Length}(c) + i \text{Torsion}(c) = -2 (r u + s v) \pmod{2\pi i} \quad (3.25)$$

where $\begin{pmatrix} p & r \\ q & s \end{pmatrix} \in SL(2, \mathbb{Z})$. We have

$$\text{Im}(u \bar{v}) = -\frac{\pi}{2} \text{Length}(c) \quad (3.26)$$

According to Refs. 44, 55, we have for the hyperbolic (p, q) -Dehn surgered manifold $\mathcal{M}_{(p,q)}$ as

$$\begin{aligned} & (\text{Vol}(\mathcal{M}_{(p,q)}) + i \text{CS}(\mathcal{M}_{(p,q)})) - (\text{Vol}(\mathcal{M}) + i \text{CS}(\mathcal{M})) \\ &= -\frac{i}{4} (\Phi_{\mathcal{M}}(u) - 4 u v) - \frac{\pi}{2} (\text{Length}(c) + i \text{Torsion}(c)) \end{aligned} \quad (3.27)$$

which follows from


$$Z_{\gamma}(\mathcal{M}_{(p,q)}) \sim \int du e^{\frac{1}{2i\gamma}(\frac{p}{q}u^2 + \frac{2(\pi+\gamma)i}{q}u)} Z_{\gamma}(\mathcal{M}_u) \quad (3.28)$$

4. E

We explain our constructions of the quantum invariant by taking some concrete examples of cusped hyperbolic 3-manifolds.

4.1 Figure-Eight knot 4_1

We set $\mathcal{K}$ as the Figure-Eight knot 4_1 which is depicted as



It is well known that the complement of the figure eight knot, $\mathcal{M} = S^3 \setminus \mathcal{K}$, is given by two ideal tetrahedra, and the triangulation induces the quantum invariant as

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_1, p_2 | S | p_3, p_4 \rangle \langle p_4, p_3 | S^{-1} | p_2, p_1 \rangle \quad (4.1)$$


Modulus of two tetrahedra are given by $w = e^{p_4 - p_2}$ and $z = e^{p_1 - p_3}$. The developing map is drawn in Fig. 3, and the meridian is read to be

$$\frac{w}{z} = e^{-2u}$$

which shows that the condition $\delta_C(\mathbf{p}; u)$ is

$$p_4 - p_2 - (p_1 - p_3) = -2u$$

The complete hyperbolic structure is realized when $u = 0$.



F 3. Developing map of the complement of the figure-eight knot. The gray filled triangle corresponds to top vertex of the tetrahedron (central vertex of circle in (3.1)) with modulus w in projection of (3.1). A curve denotes a meridian of cusp.

We then obtain

$$Z_\gamma(\mathcal{M}_u) = \frac{1}{4\pi\gamma} \int dx \frac{\Phi_\gamma(x + i\pi + i\gamma)}{\Phi_\gamma(-x - 2u - i\pi - i\gamma)} e^{\frac{-1}{2i\gamma} 4u(u+x)} \quad (4.2)$$

In a limit $\gamma \rightarrow 0$, we have

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) &\sim \int dx \exp\left(\frac{1}{2i\gamma} \left(\text{Li}_2(e^x) - \text{Li}_2(e^{-x-2u}) - 4u(u+x)\right)\right) \\ &= \int dx \exp\left(\frac{1}{2i\gamma} V_{\mathcal{M}}(e^x; e^u)\right) \end{aligned} \quad (4.3)$$

where the potential function is set to be

$$V_{\mathcal{M}}(x; m) = \text{Li}_2(x) - \text{Li}_2\left(\frac{1}{xm^2}\right) - 4 \log m \log(xm) \quad (4.4)$$

To evaluate the integral (4.3), we may apply the saddle point method, and a condition (3.11) reduces to

$$-\frac{x}{m^2(1-x)(1-m^2x)} = 1 \quad (4.5)$$

From (3.12) the longitude is given by

$$\frac{1}{m^2x(m^2x-1)} = \ell \quad (4.6)$$

Completeness condition $m = 1$ gives $x = \frac{1 \pm \sqrt{3}i}{2}$ from (4.5). Substituting this solution for the potential function (4.4), the imaginary part coincides with the hyperbolic volume of the figure-eight knot $\text{Vol}(S^3 \setminus 4_1) = 2.02988 \dots$.

For a deformed manifold $\mathcal{M}_u$, we get an algebraic equation of ℓ and m by eliminating x from (4.5) and (4.6) as

$$A(\ell, m) = 0 \quad (4.7)$$

Here $A(\ell, m)$ is the A-polynomial for the figure-eight knot;

$$A(\ell, m) = -m^4 + \ell(1 - m^2 - 2m^4 - m^6 + m^8) - \ell^2 m^4 \quad (4.8)$$

which can be expressed as in the Newton polygon as follows;

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & -1 & 0 \\ -1 & -2 & -1 \\ 0 & -1 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$


We note that a set of equations (4.5) and (4.6) gives

$$2v = -2\pi i + 4\sqrt{3}iu + \frac{16i}{\sqrt{3}}u^3 + \frac{368i}{15\sqrt{3}}u^5 + \frac{2848i}{45\sqrt{3}}u^7 + \dots$$

which coincides with a result in Ref. 44 under (3.15).

4.2 5_2

We set $\mathcal{K}$ as the knot 5_2 depicted as



The knot complement $\mathcal{M} = S^3 \setminus \mathcal{K}$ is constructed from three tetrahedra (see, *e.g.*, Ref. 49) and we obtain the invariant as


$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_1, p_5 | S^{-1} | p_4, p_3 \rangle \langle p_2, p_4 | S^{-1} | p_6, p_5 \rangle \langle p_3, p_6 | S^{-1} | p_1, p_2 \rangle \quad (4.9)$$

Modulus of three tetrahedra are respectively $z_1 = e^{p_3 - p_5}$, $z_2 = e^{p_5 - p_4}$, and $z_3 = e^{p_2 - p_6}$. The developing map is depicted in Fig. 4. The meridian is read to be $\frac{z_3}{z_2}$, and the condition $\delta_C(\mathbf{p}; u)$ is

$$p_5 - p_4 + p_6 - p_2 = 2u$$

We then have

$$Z_\gamma(\mathcal{M}_u) = \frac{1}{(4\pi\gamma)^{3/2}} \iint dx dy e^{\frac{1}{2i\gamma} \left(\frac{\pi^2 + \gamma^2}{2} + \frac{3}{2}\pi\gamma + (2u - y)(y - x) \right)} \times \frac{1}{\Phi_\gamma(-x + y - 2u - i\pi - i\gamma) \Phi_\gamma(-y - 2u - i\pi - i\gamma) \Phi_\gamma(-y - 2u - i\pi - i\gamma)} \quad (4.10)$$



F 4. Developing map of the complement of 5_2 . Gray filled triangle denotes top vertex of the tetrahedron with modulus z_1 . Meridian is denoted by a gray curve.

In a classical limit $\gamma \rightarrow 0$, we obtain

$$Z_\gamma(\mathcal{M}_u) \sim \iint dx dy \exp\left(\frac{1}{2i\gamma} V_{\mathcal{M}}(e^x, e^y; e^u)\right) \quad (4.11)$$

where the Neumann–Zagier potential function is

$$V_{\mathcal{M}}(x, y; m) = \frac{\pi^2}{2} - \text{Li}_2\left(\frac{y}{x m^2}\right) - \text{Li}_2\left(\frac{1}{y m^2}\right) - \text{Li}_2\left(\frac{1}{y}\right) + \log(y/x) \log(m^2/y) \quad (4.12)$$

The integral is evaluated by the saddle point method, and saddle point conditions (3.11) which denote hyperbolic consistency conditions reduce to

$$\begin{aligned} m^2 x - y &= x y \\ (m^2 y - 1)(y - 1) &= m^2 (m^2 x - y) \end{aligned} \quad (4.13)$$

and the identity for the longitude (3.12) is given as

$$m^4 y^2 = (m^2 x - y)(m^2 y - 1) \ell \quad (4.14)$$

In the case of complete structure $\mathcal{M}$, i.e., $u = 0$, there exists a solution of (4.13), $(x, y) = (-0.877439 + 0.744862i, 0.78492 + 1.30714i)$, such that the imaginary part of the potential function

$$\text{Im } V_{\mathcal{M}}(x, y; 1) = -D\left(\frac{y}{x}\right) - 2D\left(\frac{1}{y}\right)$$

gives the hyperbolic volume of $\mathcal{M}$; $\text{Vol}(\mathcal{M}) = 2.82812 \dots$.

To get the A-polynomial, we eliminate variables x and y from a set of equations (4.13) and (4.14). After some algebra we obtain $A_{\mathcal{K}}(\ell, m) = 0$, where the function $A_{\mathcal{K}}(\ell, m)$ coincides with the A-polynomial for 5_2 given by the following Newton polygon;

$$\begin{pmatrix} -1 & 1 & & & & \\ & -2 & & & & \\ & -2 & -1 & & & \\ & & 1 & & & \\ & 1 & & & & \\ & -1 & -2 & & & \\ & & -2 & & & \\ & & 1 & -1 & & \end{pmatrix}$$

4.3 Pretzel knot $(-2, 3, 7)$

Let $\mathcal{K}$ be the $(-2, 3, 7)$ Pretzel knot depicted as



and we set $\mathcal{M}$ as the complement of $\mathcal{K}$. The Pretzel knot $(-2, 3, 7)$ has the same hyperbolic volume with 5_2 , $\text{Vol}(\mathcal{M}) = 2.82812 \dots$, but the triangulations of the complement give the following partition function;

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_4, p_7 | S | p_6, p_1 \rangle \langle p_5, p_8 | S | p_7, p_5 \rangle \\ \times \langle p_1, p_6 | S | p_8, p_2 \rangle \langle p_3, p_2 | S | p_4, p_3 \rangle \quad (4.15)$$

Note that this triangulation differs from the canonical triangulation in Ref. 52. The developing map is given in Fig. 5. Here we set the modulus of four ideal tetrahedra as

$$\begin{aligned} z_1 &= e^{p_1 - p_7} & z_2 &= e^{p_5 - p_8} \\ z_3 &= e^{p_2 - p_6} & z_4 &= e^{p_3 - p_2} \end{aligned}$$

Then the meridian is read as


$$p_2 - p_6 - p_1 + p_7 = -2u$$

In the classical limit of the quantum invariant $Z_\gamma(\mathcal{M}_u)$ we obtain the potential function after some change of variables as

$$V_{\mathcal{M}}(x, y, z; m) = -\frac{2\pi^2}{3} + \text{Li}_2\left(\frac{1}{z}\right) + \text{Li}_2\left(\frac{1}{xyzm^4}\right) + \text{Li}_2\left(\frac{1}{zm^2}\right) + \text{Li}_2\left(\frac{m^2}{x}\right) \\ + 3\left(\log(m^2)\right)^2 + \log\left(\frac{y^5}{x^2}\right)\log(m^2) + (\log x)^2 + (\log y)^2 \quad (4.16)$$

The hyperbolic consistency conditions (3.11) give

$$\begin{aligned} (xyzm^4 - 1)(x - m^2) &= yzm^8 \\ ym^6(xyzm^4 - 1) &= xz \\ (z - 1)(xyzm^4 - 1)(m^2z - 1) &= xyz^3m^6 \end{aligned} \quad (4.17)$$



F 5. Developing map of complement of the Pretzel knot $(-2, 3, 7)$. Horosphere of the top vertex of the tetrahedron with modulus z_1 is depicted by gray triangle. Gray curve denotes a meridian.

and the longitude (3.12) is defined by

$$\ell = \frac{(xyzm^4 - 1)^2 (m^2z - 1)^2 m^2 y^3}{(x - m^2) x^3 z^3} \quad (4.18)$$

In the complete hyperbolic structure $m = 1$, we have a solution of (4.17),

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0.337641 - 0.56228i \\ 0.122561 + 0.744862i \\ 0.618504 - 0.410401i \end{pmatrix}$$

s.t. we recover the hyperbolic volume of $\mathcal{K}$;

$$\text{Im } V_M(x, y, z; m = 1) = 2D(1/z) + D\left(\frac{1}{xyz}\right) + D(1/x) = 2.82812 \dots$$

The A-polynomial is computed by eliminating (x, y, z) from (4.17) and (4.18), and we obtain $A_K(\ell, m) = 0$, where the A-polynomial for the $(-2, 3, 7)$ Pretzel knot is

$$A_K(\ell, m) = -1 + (m^{16} - 2m^{18} + m^{20})\ell + (2m^{36} + m^{38})\ell^2 - \ell^4(m^{72} + 2m^{74}) - \ell^5(m^{90} - 2m^{92} + m^{94}) + m^{110}\ell^6 \quad (4.19)$$

4.4 Once-Punctured Torus Bundles over the Circle

One of benefits of the quantum invariant $Z_\gamma(\mathcal{M})$ is that we can compute it explicitly for a once-punctured torus bundle over S^1 . A once-punctured torus bundle over S^1 , which we denote $\mathcal{M}(\varphi)$, is described by $F \times [0, 1]/(x, 0) \sim (\varphi(x), 1)$ where a monodromy matrix $\varphi \in SL(2, \mathbb{Z})$ is a homeomorphism from the punctured torus $F = \mathbb{T}^2 \setminus \{0\}$ to itself. Thurston's hyperbolization theorem indicates that $\mathcal{M}(\varphi)$ admits a complete hyperbolic metric with a finite volume when φ has 2 distinct real eigenvalues. In this case, the monodromy matrix φ can be written up to conjugation as

$$\varphi = L^{s_1} R^{t_1} \cdots L^{s_n} R^{t_n} \quad (4.20)$$

where

$$L = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad R = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$$

with $n > 0$, and s_j and t_j are positive integers. Note that the complement of the figure-eight knot studied in Sec. 4.1 corresponds to $\varphi = L R = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$.

It is known that we can triangulate the manifold M_φ with $\sum_{k=1}^n (s_k + t_k)$ ideal tetrahedra [15], and we have the quantum invariant as

$$Z_\gamma(\mathcal{M}_u(\varphi)) = \iiint_{\mathbb{R}} d\mathbf{a} d\mathbf{b} d\mathbf{c} d\mathbf{d} \delta_C(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}; u) \delta_G(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}) \times \prod_{k=1}^n \left[\prod_{i=0}^{s_k-1} \langle d_{k,i}, c_{k,i+1} | S^{-1} | d_{k,i+1}, c_{k,i} \rangle \prod_{j=0}^{t_k-1} \langle b_{k,j}, a_{k,j+1} | S | b_{k,j+1}, a_{k,j} \rangle \right] \quad (4.21)$$

where gluing condition $\delta_G(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d})$ means

$$\begin{cases} b_{k,0} = c_{k+1,0} \\ a_{k,0} = d_{k+1,0} \end{cases} \quad \text{for } k = 1, 2, \dots, n-1 \quad \begin{cases} b_{n,0} = c_{1,0} \\ a_{n,0} = d_{1,0} \end{cases}$$

$$\begin{cases} b_{k,t_k} = c_{k,s_k} \\ a_{k,t_k} = d_{k,s_k} \end{cases} \quad \text{for } k = 1, 2, \dots, n$$

When we set the modulus of each oriented tetrahedra as

$$Z_{k,j} = e^{a_{k,j} - a_{k,j+1}} \quad w_{k,j} = e^{c_{k,j} - c_{k,j+1}}$$

the developing map is drawn schematically as Fig. 6. We can then read a condition $\delta_C(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}; u)$ for meridian as

$$u = \frac{1}{2} \sum_{k=1}^n (c_{k,0} - c_{k,s_k} - a_{k,0} + a_{k,t_k}) \quad (4.22)$$

With these conditions, we have the quantum invariant $Z_\gamma(\mathcal{M}_u(\varphi))$, and the Neumann–Zagier potential function can be given by taking a classical limit $\gamma \rightarrow 0$. As far as we know, the A-polynomial-type invariant for the once-punctured torus bundle over S^1 is not studied, but we can obtain such polynomials from the Neumann–Zagier function.

Below we give a few examples for concreteness.

4.4.1 $L^2 R$

We set $\varphi = L^2 R = \begin{pmatrix} 3 & 2 \\ 1 & 1 \end{pmatrix}$. The hyperbolic Dehn surgery of this manifold is studied in Ref. 4. The invariant (4.21) is rewritten as

$$\begin{aligned} Z_\gamma(\mathcal{M}_u(L^2 R)) \\ = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_1, p_5 | S^{-1} | p_6, p_3 \rangle \langle p_6, p_4 | S^{-1} | p_2, p_5 \rangle \langle p_3, p_2 | S | p_4, p_1 \rangle \end{aligned} \quad (4.23)$$

When we set the modulus of tetrahedra as $w_0 = e^{p_3 - p_5}$, $w_1 = e^{p_5 - p_4}$, and $z_0 = e^{p_1 - p_2}$, the developing map $\mathcal{M}(L^2 R)$ can be depicted as Fig. 7. Then the meridian is read as


$$p_3 - p_4 - p_1 + p_2 = 2u \quad (4.24)$$

In the classical limit, we have

$$Z_\gamma(\mathcal{M}_u(L^2 R)) \sim \iint dx dy \exp\left(\frac{1}{2i\gamma} V_{\mathcal{M}(L^2 R)}(e^x, e^y; e^u)\right) \quad (4.25)$$

where the potential function is computed as

$$\begin{aligned} V_{\mathcal{M}(L^2 R)}(x, y; m) = \text{Li}_2\left(\frac{1}{m^2 x}\right) - \text{Li}_2\left(\frac{1}{m^2 x y^2}\right) - \text{Li}_2(m^2 x^2 y^2) \\ - \log(m^2) \log(m^2 x^2 y^4) - 2(\log(xy))^2 + \frac{\pi^2}{6} \end{aligned} \quad (4.26)$$




F 6. Schematic developing map for $\mathcal{M}(\varphi)$. Top vertex of tetrahedron with modulus $z_{1,0}$ is filled gray. Gray straight line denotes meridian.

The saddle point conditions (3.11), which correspond to the hyperbolic consistency condition, reduce to

$$\frac{(-1 + m^2 x) (-1 + m^2 x^2 y^2)^2}{m^4 x^4 y^2 (-1 + m^2 x y^2)} = 1$$

$$\frac{(-1 + m^2 x^2 y^2)}{m^2 x (-1 + m^2 x y^2)} = 1$$
(4.27)



F 7. Developing map of $\mathcal{M}(L^2 R)$. Gray filled triangle is top vertex of the tetrahedron with modulus z_0 . Gray curve is meridian.

and, as (3.12), the longitude is defined by

$$\ell = -\frac{(-1 + m^2 x)(-1 + m^2 x^2 y^2)}{m^4 x^2 y^2 (-1 + m^2 x y^2)} \quad (4.28)$$

In the case of complete case $m = 1$, we find that $(x, y^2) = \left(\frac{1 \pm \sqrt{7}i}{4}, \frac{-1 \pm \sqrt{7}i}{2}\right)$ solves the consistency condition (4.27). Among these, we can check numerically that the largest value of the imaginary part of the potential function at the saddle points

$$\text{Im } V_{\mathcal{M}(L^2 R)}(x, y; m = 1) = D\left(\frac{1}{x}\right) - D\left(\frac{1}{x y^2}\right) - D(x^2 y^2)$$

coincides with the hyperbolic volume $\text{Vol}(\mathcal{M}(L^2 R)) = 2.66674 \dots$.

The A-polynomial is now given by eliminating x and y from the set of equations, (4.27) and (4.28), and we obtain the algebraic curve $A_{\mathcal{M}(L^2 R)}(\ell, m) = 0$ whose Newton polygon is given by

$$\begin{pmatrix} 0 & -1 & 0 \\ 0 & 2 & 1 \\ 1 & 2 & 0 \\ 0 & -1 & 0 \end{pmatrix}$$

4.4.2 LR^3

We take another example $\varphi = LR^3 = \begin{pmatrix} 4 & 1 \\ 3 & 1 \end{pmatrix}$. In this case, the invariant (4.21) becomes


$$Z_\gamma(\mathcal{M}_u(LR^3)) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_1, p_4 | S | p_3, p_2 \rangle \langle p_3, p_6 | S | p_5, p_4 \rangle \\ \times \langle p_5, p_8 | S | p_7, p_6 \rangle \langle p_2, p_7 | S^{-1} | p_8, p_1 \rangle \quad (4.29)$$

We set the modulus of each tetrahedron as

$$\begin{aligned} z_0 &= e^{p_2 - p_4} & z_1 &= e^{p_4 - p_6} \\ z_2 &= e^{p_6 - p_8} & w_0 &= e^{p_1 - p_7} \end{aligned}$$

The developing map is depicted in Fig. 8, and the meridian is read as $\frac{w_0}{z_0 z_1 z_2}$. We thus have a condition of $\delta_C(\mathbf{p}; u)$ as

$$p_1 - p_7 - p_2 + p_8 = 2u \quad (4.30)$$



F 8. Developing map of $\mathcal{M}(LR^3)$.

In the classical limit, we obtain

$$Z_\gamma(\mathcal{M}_u(LR^3)) \sim \iiint dx dy dz \exp\left(\frac{1}{2i\gamma} V_{\mathcal{M}(LR^3)}(e^x, e^y, e^z; e^u)\right) \quad (4.31)$$

where the potential function is computed as

$$\begin{aligned} V_{\mathcal{M}(LR^3)}(x, y, z; m) &= -\text{Li}_2\left(\frac{1}{m^4 y}\right) + \text{Li}_2\left(\frac{1}{m^2 x y z^2}\right) + \text{Li}_2(m^2 x^2 z) \\ &\quad + \text{Li}_2\left(\frac{m^2 y^2 z}{x}\right) - (\log(m^2))^2 + 2 \log\left(\frac{x}{m^2}\right) \log\left(\frac{x}{y}\right) \\ &\quad + 2 \log y \log(y z) + 2 \log z \log(x z m^2) - \frac{\pi^2}{3} \end{aligned} \quad (4.32)$$

In the case of complete case $m = 1$, we can check numerically that among algebraic solutions of (3.11) the maximum of the imaginary part of $V_{\mathcal{M}(LR^3)}(x, y, z; m = 1)$,

$$\operatorname{Im} V_{\mathcal{M}(LR^3)}(x, y, z; 1) = -D\left(\frac{1}{y}\right) + D\left(\frac{1}{xy z^2}\right) + D(x^2 z) + D\left(\frac{y^2 z}{x}\right)$$

coincides with the hyperbolic volume $\operatorname{Vol}(\mathcal{M}(LR^3)) = 2.98912 \dots$. See Ref. 16 where algebraic solutions of consistency equations are investigated in detail.

Correspondingly we obtain the A-polynomial for $\mathcal{M}(LR^3)$ by eliminating x , y , and z from a set of equations. Explicitly the polynomial $A_{\mathcal{M}(LR^3)}(\ell, m)$ is given in the form of Newton polygon as follows;

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -3 & 0 & 0 \\ -1 & -2 & -3 & -1 & -2 & 0 \\ 0 & -3 & -2 & 2 & 3 & 0 \\ 0 & 2 & 1 & 3 & 2 & 1 \\ 0 & 0 & 3 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \end{pmatrix}$$

5. C

D

We have constructed quantum invariant $Z_\gamma(\mathcal{M})$ for cusped hyperbolic 3-manifolds $\mathcal{M}$ by assigning the Faddeev quantum dilogarithm function to oriented ideal tetrahedra. Once the triangulation of cusped 3-manifold $\mathcal{M}$ is given, it is rather straightforward to define the invariant $Z_\gamma(\mathcal{M})$ in an integral form. In the classical limit, the Faddeev integral reduces to the dilogarithm function, which denotes the hyperbolic volume of ideal tetrahedron. Remarkable is that the saddle point conditions coincide with the hyperbolic consistency conditions around edges [19]. We have discussed as a variant of the volume conjecture that the invariant $Z_\gamma(\mathcal{M})$, which can be regarded as a generalization of the Kashaev invariant (specific value of the colored Jones polynomial), is dominated by the hyperbolic volume in the classical limit $\gamma \rightarrow 0$.

We have shown that the invariant can be defined for a one-parameter deformation of manifold $\mathcal{M}_u$ (not complete), and that the Neumann–Zagier potential function can be given by taking a classical limit $\gamma \rightarrow 0$. Correspondingly the A-polynomial can be computed from the potential function (3.14). This supports the generalized volume conjecture. We have demonstrated by taking examples that our method recovers previously known A-polynomial when $\mathcal{M}$ is a complement of hyperbolic knots. We have further applied our method in the case of the once-punctured torus bundle over the circle. It seems that the A-polynomial-type invariant is not known for this

3-manifold, and it will be interesting to study a relationship with the boundary slope. To conclude, our results indicate that the S -operator (2.13) denotes the quantum Bloch invariant for oriented ideal tetrahedron.

The A-polynomial may give an interesting insight for mathematical physics. For example, the Mahler measure of the A-polynomial is expected to coincide with the hyperbolic volume of knot. On the other hand, the Mahler measure of the determinant of the Kasteleyn matrix has appeared as the free energy of the dimer problem. From the viewpoint of our combinatorial construction of the quantum invariant, the quantum invariants based on oriented ideal triangulation may be interpreted as a matching of in-states and out-states, and it might be interesting to investigate a Kasteleyn matrix interpretation of the A-polynomial.

A

We have used computer programs, SnapPea [52], Knotscape [24], and Snap [11], in studying triangulation of manifolds. We have also used Mathematica and Pari/GP. Pictures of knots in this paper are drawn using KnotPlot [47]. This work is supported in part by Grant-in-Aid for Young Scientists from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

A. M E

We shall study the quantum invariant for other cusped manifolds. We give


- the quantum invariant $Z_\gamma(\mathcal{M})$ in terms of the S -operators,
- developing map when the number of the ideal tetrahedron is less than four,
- the condition $\delta_C(\mathbf{p}; u)$, where u is a deformation parameter from the completeness $u = 0$,
- the Neumann–Zagier potential function $V_{\mathcal{M}}(\mathbf{x}; m)$, which follows from $Z_\gamma(\mathcal{M})$ by taking the classical limit $\gamma \rightarrow 0$,
- a solution of the saddle point equations in the complete case $u = 0$, which is checked that the hyperbolic volume coincides with a maximal value of the imaginary part of the potential function at this saddle point,
- the A-polynomial, which is given from the potential function $V_{\mathcal{M}}(\mathbf{x}; m)$ by eliminating parameters $\mathbf{x}$.

In the first part A.1, we collect results for hyperbolic knots up to 7 crossings. Though the A-polynomial is given in Ref. 9, we give an explicit form for self-completeness. In the second

part A.2, we consider *simple* hyperbolic knots K_{x_y} from Ref. 6. Pictures of knot may be complicated, but the triangulation of the complement is relatively simple in these cases. We note that, though the number of the ideal tetrahedra of the complement of knot K_{x_y} is x in the case of the canonical triangulation [52], our triangulations are different.

A.1 Complement of Knots up to 7 Crossings

A.1.1 6_1




$$\text{Vol}(S^3 \setminus \mathcal{K}) = 3.16396 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_3, p_1 | S^{-1} | p_2, p_4 \rangle \langle p_6, p_4 | S | p_5, p_8 \rangle \\ \times \langle p_7, p_5 | S^{-1} | p_6, p_1 \rangle \langle p_8, p_2 | S | p_7, p_3 \rangle$$

- Developing map (Fig. 9)



F 9. Developing map of the knot complement of 6_1 . Gray triangle is a horosphere of the top vertex of the tetrahedron with modulus z_1 , where we have set $z_1 = e^{p_4 - p_1}$, $z_2 = e^{p_8 - p_4}$, $z_3 = e^{p_1 - p_5}$, and $z_4 = e^{p_3 - p_2}$.

- Condition $\delta_C(\mathbf{p}; u)$

$$p_4 - p_1 - p_3 + p_2 = 2u$$

- Potential function

$$V_{\mathcal{M}}(x, y, z; m) = \operatorname{Li}_2\left(\frac{zm^2}{xy}\right) + \operatorname{Li}_2(y) - \operatorname{Li}_2\left(\frac{m^2}{y}\right) - \operatorname{Li}_2\left(\frac{yz}{m^2}\right) \\ + \log(m^2) \log\left(\frac{yz^2}{x}\right) - \log z \log(xy)$$

- Hyperbolic volume

$$\operatorname{Vol}(\mathcal{M}) = \operatorname{Im} V_{\mathcal{M}}(x, y, z; m = 1) = D\left(\frac{z}{xy}\right) + D(y) - D\left(\frac{1}{y}\right) - D(yz)$$


where

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.851808 + 0.911292i \\ 0.278726 + 0.48342i \\ -1.50411 - 1.22685i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 3 & -1 & 0 \\ 0 & 2 & 1 & 0 & 0 \\ 0 & -3 & -3 & 0 & 0 \\ -1 & -3 & -6 & -3 & -1 \\ 0 & 0 & -3 & -3 & 0 \\ 0 & 0 & 1 & 2 & 0 \\ 0 & -1 & 3 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \end{pmatrix}$$

A.1.2 6_2



$$\operatorname{Vol}(S^3 \setminus \mathcal{K}) = 4.40083 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_8, p_6 | S^{-1} | p_1, p_2 \rangle \langle p_3, p_7 | S^{-1} | p_8, p_4 \rangle \\ \times \langle p_4, p_5 | S | p_3, p_9 \rangle \langle p_2, p_{10} | S | p_6, p_5 \rangle \langle p_1, p_9 | S^{-1} | p_{10}, p_7 \rangle$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_2 - p_6 - p_5 + p_{10} = -2u$$

- Potential function

$$V_{\mathcal{M}}(w, x, y, z; m) = \text{Li}_2\left(\frac{z}{m^2}\right) + \text{Li}_2\left(\frac{m^4}{yw}\right) - \text{Li}_2\left(\frac{yw}{m^2}\right) - \text{Li}_2\left(\frac{m^2}{xy}\right) - \text{Li}_2\left(\frac{yz}{m^2}\right) \\ + \log x \log z - \log(yw) \log(yz) + \log(m^2) \log\left(\frac{y^2 z}{w}\right)$$

- Hyperbolic volume


$$\text{Im } V_{\mathcal{M}}(w, x, y, z; 1) = D(z) + D\left(\frac{1}{yw}\right) - D(yw) - D\left(\frac{1}{xy}\right) - D(yz)$$

with

$$\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.455697 + 1.20015 i \\ -0.964913 - 0.621896 i \\ -0.418784 - 0.219165 i \\ 0.0904327 + 1.60288 i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -2 & 1 & 0 & 0 & 0 \\ -1 & 1 & -3 & 0 & 0 & 0 \\ 0 & 2 & 1 & 0 & 0 & 0 \\ 0 & -5 & 5 & 0 & 0 & 0 \\ 0 & -5 & 3 & -3 & 0 & 0 \\ 0 & 3 & -12 & 8 & 0 & 0 \\ 0 & 0 & -13 & 3 & 0 & 0 \\ 0 & 0 & 3 & -13 & 0 & 0 \\ 0 & 0 & 8 & -12 & 3 & 0 \\ 0 & 0 & -3 & 3 & -5 & 0 \\ 0 & 0 & 0 & 5 & -5 & 0 \\ 0 & 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & -3 & 1 & -1 \\ 0 & 0 & 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}$$

A.1.3 6_3 

$$\text{Vol}(S^3 \setminus \mathcal{K}) = 5.69302 \dots$$

- Quantum invariant

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \, \delta_C(\mathbf{p}; u) & \langle p_3, p_2 | S | p_1, p_9 \rangle \langle p_7, p_9 | S^{-1} | p_5, p_{10} \rangle \langle p_{10}, p_4 | S^{-1} | p_6, p_8 \rangle \\ & \times \langle p_8, p_1 | S^{-1} | p_4, p_{11} \rangle \langle p_{12}, p_{11} | S^{-1} | p_2, p_3 \rangle \langle p_6, p_5 | S | p_{12}, p_7 \rangle \quad (\text{A.1}) \end{aligned}$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_{10} - p_9 - p_7 + p_5 = -2u$$

- Potential function

$$\begin{aligned} V_{\mathcal{M}}(v, w, x, y, z; m) \\ = \text{Li}_2\left(\frac{m^2 y}{x}\right) + \text{Li}_2\left(\frac{v}{w}\right) - \text{Li}_2\left(\frac{m^2 w}{v}\right) - \text{Li}_2\left(\frac{xz}{v}\right) - \text{Li}_2\left(\frac{x^2 z}{v}\right) - \text{Li}_2\left(\frac{v}{xyz}\right) \\ + \frac{\pi^2}{3} + \log(m^2) \log\left(\frac{y}{w}\right) + \log x \log\left(\frac{v}{yz^2}\right) + \log\left(\frac{w}{yz}\right) \log\left(\frac{z}{v}\right) \end{aligned}$$

- Hyperbolic volume

$$\text{Im } V_{\mathcal{M}}(v, w, x, y, z; 1) = D\left(\frac{y}{x}\right) + D\left(\frac{v}{w}\right) - D\left(\frac{w}{v}\right) - D\left(\frac{xz}{v}\right) - D\left(\frac{x^2 z}{v}\right) - D\left(\frac{v}{xyz}\right)$$


with

$$\begin{pmatrix} v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0.0739495 + 0.558752 i \\ 0.732786 + 0.381252 i \\ 1.0 \\ 0.108378 + 0.818891 i \\ 0.415113 + 0.381252 i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -5 & 1 & 0 & 0 \\ 0 & 0 & -4 & 3 & -4 & 0 & 0 \\ 0 & 0 & 4 & 9 & 4 & 0 & 0 \\ 0 & 2 & 2 & -2 & 2 & 2 & 0 \\ 0 & -5 & -6 & -21 & -6 & -5 & 0 \\ 0 & 1 & 2 & 8 & 2 & 1 & 0 \\ 1 & 10 & 17 & 34 & 17 & 10 & 1 \\ 0 & 1 & 2 & 8 & 2 & 1 & 0 \\ 0 & -5 & -6 & -21 & -6 & -5 & 0 \\ 0 & 2 & 2 & -2 & 2 & 2 & 0 \\ 0 & 0 & 4 & 9 & 4 & 0 & 0 \\ 0 & 0 & -4 & 3 & -4 & 0 & 0 \\ 0 & 0 & 1 & -5 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

A.1.4 7_2



$$\text{Vol}(S^3 \setminus \mathcal{K}) = 3.33174 \dots$$

- Quantum invariant


$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \, \delta_C(\mathbf{p}; u) \langle p_1, p_4 | S | p_2, p_3 \rangle \langle p_5, p_2 | S | p_8, p_1 \rangle \\ \times \langle p_8, p_7 | S^{-1} | p_7, p_6 \rangle \langle p_6, p_3 | S | p_4, p_5 \rangle$$

- Developing map (Fig. 10)
- Condition $\delta_C(\mathbf{p}; u)$

$$p_3 - p_4 - p_1 + p_2 = -2u$$

- Potential function

$$V_{\mathcal{M}}(x, y, z; m) = \text{Li}_2\left(\frac{1}{x}\right) + \text{Li}_2\left(\frac{m^2}{x}\right) + \text{Li}_2\left(z m^2\right) - \text{Li}_2\left(\frac{y z}{x^2}\right) - \frac{\pi^2}{3} \\ - (\log(x/y))^2 + \log z \log\left(\frac{x^3 m^2}{y^2 z}\right)$$



F 10. Developing map of the knot complement of 7_2 . The gray triangle corresponds to top vertex of the tetrahedron with modulus z_1 , where we have set modulus as $z_1 = e^{p_3 - p_4}$, $z_2 = e^{p_1 - p_2}$, $z_3 = e^{p_6 - p_7}$, and $z_4 = e^{p_5 - p_3}$.

- Hyperbolic volume


$$\text{Im } V_{\mathcal{M}}(x, y, z; 1) = 2D\left(\frac{1}{x}\right) + D(z) - D\left(\frac{yz}{x^2}\right)$$

with

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0.941819 - 1.69128i \\ 0.935538 + 0.903908i \\ 0.0581814 + 1.69128i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 1 & -2 & 1 & 0 & 0 & 0 \\ 0 & 4 & -4 & 0 & 0 & 0 \\ 0 & 3 & 2 & -2 & 0 & 0 \\ 0 & 0 & 5 & 5 & 0 & 0 \\ 0 & 0 & 6 & 1 & 1 & 0 \\ 0 & 0 & 0 & -4 & -1 & 0 \\ 0 & -1 & -4 & 0 & 0 & 0 \\ 0 & 1 & 1 & 6 & 0 & 0 \\ 0 & 0 & 5 & 5 & 0 & 0 \\ 0 & 0 & -2 & 2 & 3 & 0 \\ 0 & 0 & 0 & -4 & 4 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{pmatrix}$$

A.1.5 7_3 

$$\text{Vol}(S^3 \setminus \mathcal{K}) = 4.5921 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \, \delta_C(\mathbf{p}; u) \langle p_1, p_3 | S | p_4, p_8 \rangle \langle p_2, p_8 | S | p_5, p_1 \rangle \\ \times \langle p_5, p_9 | S^{-1} | p_6, p_7 \rangle \langle p_4, p_6 | S | p_3, p_{10} \rangle \langle p_7, p_{10} | S | p_2, p_9 \rangle$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_{10} - p_6 - p_1 + p_8 = -2u$$

- Potential function

$$V_{\mathcal{M}}(w, x, y, z; m) = \text{Li}_2(m^2 w) + \text{Li}_2(m^4 w) + \text{Li}_2\left(\frac{1}{m^2 x}\right) + \text{Li}_2\left(\frac{z}{w}\right) - \text{Li}_2\left(\frac{z}{y}\right) - \frac{\pi^2}{2} \\ + \log w \log x + \log y \log z + \log(m^2) \log(y w^2 z)$$

- Hyperbolic volume

$$\text{Im } V_{\mathcal{M}}(w, x, y, z; 1) = 2 D(w) + D\left(\frac{1}{x}\right) + D\left(\frac{z}{w}\right) - D\left(\frac{z}{y}\right)$$


with

$$\begin{pmatrix} x \\ y \\ z \\ w \end{pmatrix} = \begin{pmatrix} 0.645284 - 0.801205 \, \text{i} \\ -0.676708 + 0.260961 \, \text{i} \\ -0.87287 + 1.51178 \, \text{i} \\ 0.537981 + 1.04357 \, \text{i} \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -5 & -3 & 0 & 0 & 0 & 0 \\ 0 & -2 & 9 & 0 & 0 & 0 & 0 \\ 0 & 3 & -2 & 0 & 0 & 0 & 0 \\ 0 & -2 & -14 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 3 & 0 & 0 & 0 \\ 0 & 0 & 4 & -10 & 0 & 0 & 0 \\ 0 & 0 & -4 & 3 & 0 & 0 & 0 \\ 0 & 0 & -2 & 12 & 0 & 0 & 0 \\ 0 & 0 & -3 & -6 & -1 & 0 & 0 \\ 0 & 0 & 3 & 24 & 3 & 0 & 0 \\ 0 & 0 & -1 & -6 & -3 & 0 & 0 \\ 0 & 0 & 0 & 12 & -2 & 0 & 0 \\ 0 & 0 & 0 & 3 & -4 & 0 & 0 \\ 0 & 0 & 0 & -10 & 4 & 0 & 0 \\ 0 & 0 & 0 & 3 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -14 & -2 & 0 \\ 0 & 0 & 0 & 0 & -2 & 3 & 0 \\ 0 & 0 & 0 & 0 & 9 & -2 & 0 \\ 0 & 0 & 0 & 0 & -3 & -5 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 \end{pmatrix}$$

A.1.6 7_4



$$\text{Vol}(S^3 \setminus \mathcal{K}) = 5.13794 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_1, p_{10} | S | p_3, p_4 \rangle \langle p_9, p_2 | S | p_{12}, p_1 \rangle \langle p_4, p_{11} | S^{-1} | p_5, p_6 \rangle \\ \times \langle p_3, p_5 | S^{-1} | p_2, p_7 \rangle \langle p_{12}, p_8 | S | p_{10}, p_9 \rangle \langle p_6, p_7 | S | p_8, p_{11} \rangle$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_4 - p_{10} + p_7 - p_5 - p_1 + p_2 = -2u$$

- Potential function

$$V_{\mathcal{M}}(v, w, x, y, z; m) = \text{Li}_2\left(\frac{w}{x}\right) + \text{Li}_2\left(\frac{vw}{y}\right) - \text{Li}_2\left(\frac{vxm^2}{y}\right) \\ + \text{Li}_2\left(\frac{m^4x}{wy}\right) - \text{Li}_2(m^2z) + \text{Li}_2\left(\frac{zx}{y}\right) - \frac{\pi^2}{3} \\ + \log(v) \log\left(\frac{w}{m^2xz}\right) - \log x \log w - 2 \log(m^2) \log z + \left(\log(m^2/w)\right)^2$$

- Hyperbolic volume

$$\text{Im } V_{\mathcal{M}}(v, w, x, y, z; 1)$$


$$= D\left(\frac{w}{x}\right) + D\left(\frac{vw}{y}\right) - D\left(\frac{vx}{y}\right) + D\left(\frac{x}{wy}\right) - D(z) + D\left(\frac{zx}{y}\right)$$

with

$$\begin{pmatrix} v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -1.10278 + 0.665457i \\ -0.102785 + 0.665457i \\ 1.0 \\ -0.664742 - 0.401127i \\ -0.226699 - 1.46771i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 0 & -1 & 3 & -3 & 1 \\ 0 & 0 & 3 & -10 & 7 & 0 \\ 0 & 0 & -3 & 3 & 4 & 0 \\ 0 & 0 & -2 & 21 & -6 & 0 \\ 0 & 1 & 10 & -3 & 1 & 0 \\ 0 & -2 & 6 & -17 & 3 & 0 \\ 0 & 3 & -17 & 6 & -2 & 0 \\ 0 & 1 & -3 & 10 & 1 & 0 \\ 0 & -6 & 21 & -2 & 0 & 0 \\ 0 & 4 & 3 & -3 & 0 & 0 \\ 0 & 7 & -10 & 3 & 0 & 0 \\ 1 & -3 & 3 & -1 & 0 & 0 \end{pmatrix}$$

A.1.7 7_5 

$$\text{Vol}(S^3 \setminus \mathcal{K}) = 6.443537 \dots$$

- Quantum invariant

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \, \delta_C(\mathbf{p}; u) & \langle p_2, p_{13} | S | p_1, p_{12} \rangle \langle p_4, p_1 | S | p_3, p_{10} \rangle \\ & \times \langle p_6, p_3 | S | p_2, p_{13} \rangle \langle p_5, p_{11} | S^{-1} | p_4, p_7 \rangle \langle p_8, p_{12} | S^{-1} | p_6, p_5 \rangle \\ & \times \langle p_7, p_9 | S | p_8, p_{14} \rangle \langle p_{10}, p_{14} | S^{-1} | p_9, p_{11} \rangle \end{aligned}$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_3 + p_7 + p_{10} = p_1 + p_5 + p_{11} - 2u$$

- Potential function


$$\begin{aligned} V_{\mathcal{M}}(u, v, w, x, y, z; m) = & -\text{Li}_2\left(\frac{u}{v}\right) + \text{Li}_2(u m^2) + \text{Li}_2\left(\frac{x}{m^2}\right) \\ & + \text{Li}_2\left(\frac{m^2}{y}\right) + \text{Li}_2\left(\frac{v}{m^2 w y}\right) - \text{Li}_2\left(\frac{v}{z}\right) - \text{Li}_2\left(\frac{w x}{z}\right) - \frac{\pi^2}{6} \\ & + \log(m^2) \log\left(\frac{u w^2}{v}\right) + \log x \log y + \log u \log z \end{aligned}$$

- Hyperbolic volume

$$\begin{aligned} \text{Im } V_{\mathcal{M}}(u, v, w, x, y, z; m = 1) = & -D\left(\frac{u}{v}\right) + D(u) + D(x) \\ & + D\left(\frac{1}{y}\right) + D\left(\frac{v}{w y}\right) - D\left(\frac{v}{z}\right) - D\left(\frac{w x}{z}\right) \end{aligned}$$

with

$$\begin{pmatrix} u \\ v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0.38762 + 1.0287 i \\ -0.572726 + 0.717749 i \\ -0.259819 + 0.832925 i \\ 0.18596 + 0.689115 i \\ 0.365014 - 1.35264 i \\ -0.679246 - 0.851242 i \end{pmatrix}$$

A.1.8 7_6 

$$\text{Vol}(S^3 \setminus \mathcal{K}) = 7.08493 \dots$$

- Quantum invariant

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) & \langle p_1, p_{14} | S^{-1} | p_3, p_{13} \rangle \langle p_4, p_{13} | S^{-1} | p_1, p_2 \rangle \langle p_9, p_5 | S | p_4, p_{14} \rangle \\ & \times \langle p_3, p_2 | S | p_6, p_{15} \rangle \langle p_7, p_6 | S | p_5, p_{16} \rangle \langle p_{11}, p_{12} | S^{-1} | p_7, p_8 \rangle \\ & \times \langle p_8, p_{15} | S^{-1} | p_6, p_{10} \rangle \langle p_{10}, p_{16} | S^{-1} | p_{11}, p_{12} \rangle \end{aligned}$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_6 + p_{10} = p_5 + p_8 - 2u$$

- Potential function

$$\begin{aligned} V_M(t, u, v, w, x, y, z; m) = & -\text{Li}_2(t) - \text{Li}_2(u m^2) + \text{Li}_2\left(\frac{v}{x}\right) - \text{Li}_2\left(\frac{m^2}{x}\right) \\ & + \text{Li}_2\left(\frac{t}{y}\right) + \text{Li}_2\left(\frac{u}{wy}\right) - \text{Li}_2\left(\frac{1}{m^2 z}\right) - \text{Li}_2\left(\frac{vw}{z}\right) + \frac{\pi^2}{3} \\ & - (\log(m^2))^2 + 2 \log(m^2) \log\left(\frac{x}{yw}\right) + \log(t/v) \log(x/y) - \log u \log z \end{aligned}$$

- Hyperbolic volume

$$\begin{aligned} \text{Im } V_M(t, u, v, w, x, y, z; 1) = & -D(t) - D(u) + D\left(\frac{v}{x}\right) - D\left(\frac{1}{x}\right) \\ & + D\left(\frac{t}{y}\right) + D\left(\frac{u}{wy}\right) - D\left(\frac{1}{z}\right) - D\left(\frac{vw}{z}\right) \end{aligned}$$


with

$$\begin{pmatrix} t \\ u \\ v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0.558614 - 1.43795 i \\ -0.0892864 - 0.842785 i \\ -0.280101 + 1.13004 i \\ 0.450985 - 0.808297 i \\ 0.234736 + 0.604244 i \\ -0.20665 - 0.833705 i \\ -0.12431 + 1.17337 i \end{pmatrix}$$

- A-polynomial

0	0	1	-1	0	0	0	0	0	0
0	0	-6	7	0	0	0	0	0	0
0	-2	11	-16	1	0	0	0	0	0
0	6	2	1	-9	0	0	0	0	0
1	-5	-16	34	32	0	0	0	0	0
0	-5	-7	10	-30	2	0	0	0	0
0	16	9	-80	-68	-16	0	0	0	0
0	5	8	-9	98	41	0	0	0	0
0	-9	42	62	164	-18	1	0	0	0
0	3	11	10	-212	-78	7	0	0	0
0	0	-37	34	-266	52	19	0	0	0
0	0	8	83	196	158	-29	0	0	0
0	0	23	-44	377	-85	10	0	0	0
0	0	-16	-48	24	-237	47	3	0	0
0	0	3	47	-237	24	-48	-16	0	0
0	0	0	10	-85	377	-44	23	0	0
0	0	0	-29	158	196	83	8	0	0
0	0	0	19	52	-266	34	-37	0	0
0	0	0	7	-78	-212	10	11	3	0
0	0	0	1	-18	164	62	42	-9	0
0	0	0	0	41	98	-9	8	5	0
0	0	0	0	-16	-68	-80	9	16	0
0	0	0	0	2	-30	10	-7	-5	0
0	0	0	0	0	32	34	-16	-5	1
0	0	0	0	0	-9	1	2	6	0
0	0	0	0	0	1	-16	11	-2	0
0	0	0	0	0	0	7	-6	0	0
0	0	0	0	0	0	-1	1	0	0

A.1.9 7_7



$$\text{Vol}(S^3 \setminus \mathcal{K}) = 7.64338 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_1, p_3 | S | p_{15}, p_{16} \rangle \langle p_2, p_4 | S | p_1, p_{14} \rangle \langle p_5, p_{13} | S^{-1} | p_8, p_4 \rangle \\ \times \langle p_6, p_{15} | S^{-1} | p_3, p_2 \rangle \langle p_7, p_{16} | S^{-1} | p_5, p_6 \rangle \langle p_{10}, p_{14} | S^{-1} | p_7, p_9 \rangle \\ \times \langle p_8, p_9 | S | p_{11}, p_{12} \rangle \langle p_{11}, p_{12} | S | p_{10}, p_{13} \rangle$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_6 + p_{15} = p_2 + p_3 - 2u$$

- Potential function

$$V_{\mathcal{M}}(t, u, v, w, x, y, z; m) = \text{Li}_2(v m^2) + \text{Li}_2(t m^2 w) - \text{Li}_2\left(\frac{t m^2}{x}\right) - \text{Li}_2\left(\frac{1}{m^2 w x}\right) \\ - \text{Li}_2(v x) + \text{Li}_2(u x) + \text{Li}_2\left(\frac{1}{m^4 z}\right) - \text{Li}_2\left(\frac{u}{m^2 z}\right) \\ + \log(m^2) \log\left(\frac{v u}{m^2 w y}\right) + \log(t u) \log\left(\frac{w x}{y}\right) + \log v \log z$$

- Hyperbolic volume

$$\text{Im } V_{\mathcal{M}}(t, u, v, w, x, y, z; 1) = D(v) + D(t w) - D\left(\frac{t}{x}\right) \\ - D\left(\frac{1}{w x}\right) - D(v x) + D(u x) + D\left(\frac{1}{z}\right) - D\left(\frac{u}{z}\right)$$

with

$$\begin{pmatrix} t \\ u \\ v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.899232 + 0.400532 i \\ -0.927958 - 0.413327 i \\ 0.0287264 + 0.813859 i \\ -0.351808 - 0.720342 i \\ -0.927958 - 0.413327 i \\ -0.927958 - 0.413327 i \\ 0.0433154 - 1.22719 i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 0 & 0 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -7 & 8 & 0 & 0 & 0 \\ 0 & 0 & 0 & 15 & -19 & 1 & 0 & 0 \\ 0 & 0 & 3 & 1 & -4 & -6 & 0 & 0 \\ 0 & 0 & -18 & -30 & 59 & 11 & 0 & 0 \\ 0 & 0 & 23 & -1 & 0 & -1 & 0 & 0 \\ 0 & 3 & 27 & 41 & -123 & -8 & -2 & 0 \\ 0 & -11 & -65 & 7 & -2 & 1 & 7 & 0 \\ 0 & 4 & -19 & -29 & 130 & -28 & -7 & 0 \\ 1 & 20 & 84 & -46 & 35 & 16 & -7 & 0 \\ 0 & -7 & 16 & 35 & -46 & 84 & 20 & 1 \\ 0 & -7 & -28 & 130 & -29 & -19 & 4 & 0 \\ 0 & 7 & 1 & -2 & 7 & -65 & -11 & 0 \\ 0 & -2 & -8 & -123 & 41 & 27 & 3 & 0 \\ 0 & 0 & -1 & 0 & -1 & 23 & 0 & 0 \\ 0 & 0 & 11 & 59 & -30 & -18 & 0 & 0 \\ 0 & 0 & -6 & -4 & 1 & 3 & 0 & 0 \\ 0 & 0 & 1 & -19 & 15 & 0 & 0 & 0 \\ 0 & 0 & 0 & 8 & -7 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 1 & 0 & 0 & 0 \end{pmatrix}$$

A.2 Complement of “Simple” Hyperbolic Knots

A.2.1 $K4_4$



$$\text{Vol}(S^3 \setminus \mathcal{K}) = 3.608689 \dots$$

- Quantum invariant

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_3, p_2 | S^{-1} | p_2, p_1 \rangle \langle p_{12}, p_5 | S | p_{10}, p_3 \rangle \langle p_6, p_1 | S^{-1} | p_{11}, p_{12} \rangle \\ \times \langle p_{10}, p_9 | S | p_8, p_6 \rangle \langle p_4, p_7 | S | p_9, p_4 \rangle \langle p_{11}, p_8 | S | p_7, p_5 \rangle \end{aligned}$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_{12} - p_1 + p_5 - p_8 - p_6 + p_9 = -2u$$

- Potential function

$$\begin{aligned}
 V_{\mathcal{M}}(v, w, x, y, z; m) = & -\operatorname{Li}_2\left(\frac{x}{w}\right) + \operatorname{Li}_2\left(\frac{1}{v m^2 y}\right) - \operatorname{Li}_2\left(\frac{z}{v m^2}\right) \\
 & + \operatorname{Li}_2(m^2 x y) + \operatorname{Li}_2\left(\frac{m^2}{y z}\right) + \operatorname{Li}_2\left(\frac{w y}{z}\right) - \frac{\pi^2}{3} + \log\left(\frac{v}{z}\right) \log(m^4 x y) \\
 & + \log(x) \log\left(\frac{m^2 w^2}{x}\right) + 2 \log(w y) \log(y) + \log(m^2) \log(m^4 w^3)
 \end{aligned}$$

- Hyperbolic volume

$$\begin{aligned}
 \operatorname{Im} V_{\mathcal{M}}(v, w, x, y, z; 1) = & -D\left(\frac{x}{w}\right) + D\left(\frac{1}{v y}\right) - D\left(\frac{z}{v}\right) \\
 & + D(x y) + D\left(\frac{1}{y z}\right) + D\left(\frac{w y}{z}\right)
 \end{aligned}$$


with

$$\begin{pmatrix} v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.06796 - 1.03267 i \\ -0.597112 + 0.762045 i \\ 1.29516 + 0.539127 i \\ 0.457778 + 1.02559 i \\ -0.396648 - 0.345221 i \end{pmatrix}$$

- A-polynomial

$$\begin{aligned}
 A(\ell, m) = & 1 + (m^{30} - 2 m^{32} + m^{34}) \ell + (-m^{58} + 2 m^{60} - 10 m^{62} + 4 m^{64} - m^{66}) \ell^2 \\
 & + (-2 m^{90} + 3 m^{92} - m^{96}) \ell^3 + (m^{120} + 8 m^{122} + 6 m^{124}) \ell^4 \\
 & + (m^{150} - m^{152} - m^{154} + m^{156}) \ell^5 + (2 m^{180} - 12 m^{182} - 12 m^{186} + 2 m^{188}) \ell^6 \\
 & + (m^{212} - m^{214} - m^{216} + m^{218}) \ell^7 + (6 m^{244} + 8 m^{246} + m^{248}) \ell^8 \\
 & + (-m^{272} + 3 m^{276} - 2 m^{278}) \ell^9 + (-m^{302} + 4 m^{304} - 10 m^{406} + 2 m^{408} - m^{310}) \ell^{10} \\
 & + (m^{334} - 2 m^{336} + m^{338}) \ell^{11} + m^{368} \ell^{12}
 \end{aligned}$$

A.2.2 $K5_1$



$$\operatorname{Vol}(S^3 \setminus \mathcal{K}) = 3.4179148 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_5, p_2 | S^{-1} | p_1, p_3 \rangle \langle p_1, p_9 | S^{-1} | p_9, p_2 \rangle \langle p_4, p_3 | S^{-1} | p_5, p_7 \rangle \\ \times \langle p_{10}, p_6 | S^{-1} | p_4, p_{11} \rangle \langle p_{12}, p_7 | S^{-1} | p_6, p_{10} \rangle \langle p_{11}, p_8 | S^{-1} | p_8, p_{12} \rangle$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_{11} - p_6 - p_{10} + p_7 = -2u$$

- Potential function

$$V_M(v, w, x, y, z; m) = -\text{Li}_2\left(\frac{w}{x^2}\right) - \text{Li}_2\left(\frac{wx}{y}\right) - \text{Li}_2\left(\frac{y^2}{m^4 v z}\right) - \text{Li}_2\left(\frac{y^2}{w z}\right) \\ - \text{Li}_2\left(\frac{z}{v y}\right) - \text{Li}_2\left(\frac{m^2 z}{v y}\right) + \pi^2 + 5 \log(m^2) \log\left(\frac{y}{z}\right) \\ + 2 \log w \log\left(\frac{x y}{w}\right) + 2 \log z \log\left(\frac{y^3}{z}\right) - 2 (\log(m^2))^2 - 2 (\log x)^2 - 5 (\log y)^2$$

- Hyperbolic volume


$$\text{Im } V_M(v, w, x, y, z; 1) = -D\left(\frac{w}{x^2}\right) - D\left(\frac{wx}{y}\right) - D\left(\frac{y^2}{v z}\right) - D\left(\frac{y^2}{w z}\right) - 2 D\left(\frac{z}{v y}\right)$$

with

$$\begin{pmatrix} v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0.465534 - 0.473866 i \\ 0.693244 + 0.159750 i \\ -1.085877 - 0.175545 i \\ -0.952444 - 0.928780 i \\ -0.907927 + 0.840443 i \end{pmatrix}$$

- A-polynomial

$$A_K(\ell, m) = -1 + (-m^{32} + m^{34}) \ell + (9m^{64} - 3m^{66} + m^{68}) \ell^2 \\ + (m^{92} - 3m^{94} + 12m^{96} - 14m^{98} + 5m^{100} - m^{102}) \ell^3 \\ + (m^{124} - 7m^{126} - 18m^{128} + 5m^{130} - 2m^{132}) \ell^4 + (-m^{156} - 7m^{158} + 2m^{160} + 6m^{162}) \ell^5 \\ + (-m^{188} + 17m^{190} + 20m^{192} - 2m^{194} + m^{196}) \ell^6 \\ + (-2m^{218} + 14m^{220} - 12m^{222} + 12m^{224} - 14m^{226} + 2m^{228}) \ell^7 \\ + (-m^{250} + 2m^{252} - 20m^{254} - 17m^{256} + m^{258}) \ell^8 \\ + (-6m^{284} - 2m^{286} + 7m^{288} + m^{290}) \ell^9 + (2m^{314} - 5m^{316} + 18m^{318} + 7m^{320} - m^{322}) \ell^{10} \\ + (m^{344} - 5m^{346} + 14m^{348} - 12m^{350} + 3m^{352} - m^{354}) \ell^{11} \\ + (-m^{378} + 3m^{380} - 9m^{382}) \ell^{12} + (-m^{412} + m^{414}) \ell^{13} + \ell^{14} m^{446}$$

A.2.3 $K5_9$ or 10_{132} 

$$\text{Vol}(S^3 \setminus \mathcal{K}) = 4.05686 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_1, p_9 | S^{-1} | p_2, p_3 \rangle \langle p_4, p_2 | S^{-1} | p_9, p_5 \rangle \\ \times \langle p_6, p_7 | S^{-1} | p_7, p_8 \rangle \langle p_5, p_3 | S | p_6, p_{10} \rangle \langle p_{10}, p_8 | S | p_4, p_1 \rangle$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_2 + p_8 + p_{10} = p_1 + p_5 + p_9 - 2u$$

- Potential function

$$V_{\mathcal{M}}(w, x, y, z; m) = \text{Li}_2(w) - \text{Li}_2\left(\frac{w}{y}\right) - \text{Li}_2\left(\frac{y}{x}\right) - \text{Li}_2(z) + \text{Li}_2\left(\frac{yz}{m^2}\right) + \frac{\pi^2}{6} \\ + \left(\log(m^2)\right)^2 - \log(m^2) \log(xyw) + \log w \log\left(\frac{y}{z}\right) + 2 \log y \log(xz)$$

- Hyperbolic volume

$$\text{Im } V_{\mathcal{M}}(w, x, y, z; 1) = D(w) - D\left(\frac{w}{y}\right) - D\left(\frac{y}{x}\right) - D(z) + D(yz)$$


with

$$\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.0498076 + 0.754729 i \\ -1.54094 - 1.35872 i \\ -0.821578 - 0.131699 i \\ -0.0844626 - 0.905094 i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & -7 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 3 & 7 & -4 & -2 & 0 & 0 \\ 0 & 1 & 4 & 2 & 13 & 4 & -6 & -3 & -1 \\ 0 & -1 & -5 & -7 & -6 & 5 & 13 & 1 & 0 \\ 0 & 0 & -5 & -12 & -18 & 9 & 9 & 4 & 0 \\ 0 & 0 & 12 & 13 & -15 & -10 & -7 & -4 & 0 \\ 0 & 4 & 7 & 10 & 15 & -13 & -12 & 0 & 0 \\ 0 & -4 & -9 & -9 & 18 & 12 & 5 & 0 & 0 \\ 0 & -1 & -13 & -5 & 6 & 7 & 5 & 1 & 0 \\ 1 & 3 & 6 & -4 & -13 & -2 & -4 & -1 & 0 \\ 0 & 0 & 2 & 4 & -7 & -3 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 7 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix}$$

A.2.4 $K5_{12}$ or 8_{20}



$$\text{Vol}(S^3 \setminus \mathcal{K}) = 4.12490 \dots$$

- Quantum invariant

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \, \delta_C(\mathbf{p}; u) \langle p_8, p_2 | S | p_1, p_6 \rangle \langle p_1, p_{10} | S^{-1} | p_{10}, p_4 \rangle \\ \times \langle p_9, p_3 | S | p_2, p_9 \rangle \langle p_4, p_7 | S^{-1} | p_5, p_3 \rangle \langle p_6, p_5 | S^{-1} | p_7, p_8 \rangle \end{aligned}$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_3 - p_7 + p_6 - p_2 - p_8 + p_5 = -2u$$

- Potential function

$$V_{\mathcal{M}}(w, x, y, z; m)$$

$$\begin{aligned} &= -\operatorname{Li}_2\left(\frac{x}{m^2 w}\right) - \operatorname{Li}_2\left(\frac{m^4 w}{xy}\right) + \operatorname{Li}_2\left(\frac{x}{y}\right) - \operatorname{Li}_2\left(\frac{z}{wy}\right) + \operatorname{Li}_2\left(\frac{z}{xy}\right) + \frac{\pi^2}{6} \\ &\quad - 2\left(\log(m^2)\right)^2 - 2(\log w)^2 + (\log x)^2 \\ &\quad + \log(m^2) \log\left(\frac{xz^2}{w^5}\right) + 2\log w \log x + 2\log w \log z - 2\log x \log z \end{aligned}$$

- Hyperbolic volume

$$\operatorname{Im} V_{\mathcal{M}}(w, x, y, z; 1) = -D\left(\frac{x}{w}\right) - D\left(\frac{w}{xy}\right) + D\left(\frac{x}{y}\right) - D\left(\frac{z}{wy}\right) + D\left(\frac{z}{xy}\right)$$


with

$$\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.723387 - 0.90034i \\ -0.637406 + 0.318768i \\ -0.483596 + 0.741071i \\ 1.08906 - 0.727199i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -5 & 0 & 0 & 0 \\ 0 & -2 & 0 & -2 & 0 & 0 \\ 0 & 2 & 3 & -4 & -1 & 0 \\ 0 & 1 & -3 & 0 & -5 & -1 \\ -1 & -5 & 0 & -3 & 1 & 0 \\ 0 & -1 & -4 & 3 & 2 & 0 \\ 0 & 0 & -2 & 0 & -2 & 0 \\ 0 & 0 & 0 & -5 & 1 & 0 \\ 0 & 0 & 0 & 1 & -1 & 0 \end{pmatrix}$$

A.2.5 $K5_{13}$



$$\operatorname{Vol}(S^3 \setminus \mathcal{K}) = 4.1249 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_2, p_9 | S^{-1} | p_1, p_3 \rangle \langle p_7, p_6 | S^{-1} | p_6, p_2 \rangle \\ \times \langle p_4, p_5 | S^{-1} | p_9, p_7 \rangle \langle p_8, p_1 | S | p_4, p_{10} \rangle \langle p_{10}, p_3 | S | p_5, p_8 \rangle \quad (\text{A.2})$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_1 + p_5 + p_8 = p_7 + p_9 + p_{10} - 2u$$

- Potential function

$$V_M(w, x, y, z; m) = -\text{Li}_2(mx) + \text{Li}_2\left(\frac{x}{w}\right) - \text{Li}_2\left(\frac{1}{w^2 y}\right) - \text{Li}_2\left(\frac{z}{y}\right) \\ + \text{Li}_2\left(\frac{z}{wy}\right) + \frac{\pi^2}{6} - (\log(m^2))^2 \\ - \log(m^2) \log(wz) + \log x \log\left(\frac{z}{yw}\right) - 2 \log w \log z$$

- Hyperbolic volume

$$\text{Im } V_M(w, x, y, z; 1) = -D(x) + D\left(\frac{x}{w}\right) - D\left(\frac{1}{w^2 y}\right) - D\left(\frac{z}{y}\right) + D\left(\frac{z}{wy}\right)$$


with

$$\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -0.812447 + 0.173142 i \\ -0.0890598 - 0.727199 i \\ -1.71268 + 1.30259 i \\ 1.09977 + 1.12945 i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -10 & 1 & 3 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 8 & -14 & -11 & 8 & 0 & 0 & 0 & 0 \\ 0 & -5 & 11 & 5 & -1 & -12 & 0 & 0 & 0 & 0 \\ -1 & 5 & -10 & 14 & 29 & -12 & 1 & 0 & 0 & 0 \\ 0 & 0 & -5 & 4 & 5 & 16 & -1 & 0 & 0 & 0 \\ 0 & -1 & 9 & -13 & -19 & 25 & -7 & 2 & 0 & 0 \\ 0 & 0 & 5 & 3 & -33 & -3 & 9 & -7 & 0 & 0 \\ 0 & 0 & -7 & 9 & -3 & -33 & 3 & 5 & 0 & 0 \\ 0 & 0 & 2 & -7 & 25 & -19 & -13 & 9 & -1 & 0 \\ 0 & 0 & 0 & -1 & 16 & 5 & 4 & -5 & 0 & 0 \\ 0 & 0 & 0 & 1 & -12 & 29 & 14 & -10 & 5 & -1 \\ 0 & 0 & 0 & 0 & -12 & -1 & 5 & 11 & -5 & 0 \\ 0 & 0 & 0 & 0 & 8 & -11 & -14 & 8 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 3 & 1 & -10 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}$$

A.2.6 $K_{5_{21}}$ or 9_{46}



$$\text{Vol}(S^3 \setminus \mathcal{K}) = 4.7517 \dots$$

- Quantum invariant

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_4, p_1 | S | p_6, p_7 \rangle \langle p_2, p_9 | S^{-1} | p_1, p_8 \rangle \\ \times \langle p_7, p_8 | S^{-1} | p_3, p_4 \rangle \langle p_6, p_5 | S^{-1} | p_9, p_{10} \rangle \langle p_3, p_{10} | S^{-1} | p_5, p_2 \rangle \end{aligned}$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_1 + p_4 + p_5 = p_7 + p_9 + p_{10} - 2u$$

- Potential function

$$\begin{aligned}
 V_{\mathcal{M}}(w, x, y, z; m) = & -\operatorname{Li}_2\left(\frac{w^2 x}{m^2}\right) + \operatorname{Li}_2\left(\frac{m^2 y}{x}\right) - \operatorname{Li}_2\left(\frac{m^2}{w^2 x z}\right) \\
 & - \operatorname{Li}_2\left(\frac{z}{w}\right) - \operatorname{Li}_2\left(\frac{w y z}{m^2}\right) + \frac{\pi^2}{2} + \left(\log(m^2)\right)^2 - 2(\log w)^2 - (\log z)^2 \\
 & + \log(m^2) \log\left(\frac{y z}{x}\right) - \log w \log x - \log(y z) \log(x w)
 \end{aligned}$$

- Hyperbolic volume

$$\operatorname{Im} V_{\mathcal{M}}(w, x, y, z; 1) = -D(w^2 x) + D\left(\frac{y}{x}\right) - D\left(\frac{1}{w^2 x z}\right) - D\left(\frac{z}{w}\right) - D(w y z)$$


with

$$\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -1.21844 + 0.168108 i \\ 0.640448 - 0.637204 i \\ 1.0 \\ -0.445837 + 0.526085 i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 0 & -1 & 1 & 0 \\ 0 & 2 & 5 & -1 & 0 \\ -1 & -5 & -3 & 2 & 0 \\ 0 & 0 & -5 & -5 & 0 \\ 0 & 2 & 2 & 2 & 0 \\ 0 & -5 & -5 & 0 & 0 \\ 0 & 2 & -3 & -5 & -1 \\ 0 & -1 & 5 & 2 & 0 \\ 0 & 1 & -1 & 0 & 0 \end{pmatrix}$$

A.2.7 $K5_{22}$ or 10_{139}



$$\operatorname{Vol}(S^3 \setminus \mathcal{K}) = 4.85117 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_5, p_9 | S | p_8, p_1 \rangle \langle p_1, p_3 | S | p_2, p_9 \rangle \\ \times \langle p_6, p_7 | S | p_4, p_3 \rangle \langle p_2, p_8 | S | p_7, p_{10} \rangle \langle p_4, p_{10} | S | p_5, p_6 \rangle$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_7 + p_9 + p_{10} = p_1 + p_3 + p_8 - 2u$$

- Potential function

$$V_{\mathcal{M}}(w, x, y, z; m) = \text{Li}_2\left(\frac{1}{m^2 x}\right) + \text{Li}_2\left(\frac{m^4 w}{y}\right) + \text{Li}_2\left(\frac{w}{xz}\right) \\ + \text{Li}_2\left(\frac{z}{w}\right) + \text{Li}_2\left(\frac{wz}{y}\right) - \frac{5\pi^2}{6} + 2 \left(\log(m^2)\right)^2 \\ + \log(m^2) \log\left(\frac{x}{y^2 z}\right) + \log x \log\left(\frac{z}{y}\right) + \log w \log\left(\frac{w}{z}\right)$$

- Hyperbolic volume

$$\text{Im } V_{\mathcal{M}}(w, x, y, z; 1) = D\left(\frac{1}{x}\right) + D\left(\frac{w}{y}\right) + D\left(\frac{w}{xz}\right) + D\left(\frac{z}{w}\right) + D\left(\frac{wz}{y}\right)$$


with

$$\begin{pmatrix} w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0.660443 - 0.716885 i \\ -0.0777392 - 0.946923 i \\ -0.460355 - 1.13932 i \\ 1.0 \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 7 & 0 & 0 & 0 \\ 0 & -3 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 6 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -3 & 0 \\ 0 & 0 & 0 & 7 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

A.2.8 $K6_{10}$



$$\text{Vol}(S^3 \setminus \mathcal{K}) = 4.40083 \dots$$

- Quantum invariant

$$Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \delta_C(\mathbf{p}; u) \langle p_2, p_6 | S^{-1} | p_4, p_1 \rangle \langle p_1, p_{12} | S^{-1} | p_{10}, p_3 \rangle \langle p_3, p_8 | S | p_6, p_{11} \rangle \\ \times \langle p_7, p_9 | S^{-1} | p_9, p_8 \rangle \langle p_{10}, p_4 | S^{-1} | p_{12}, p_5 \rangle \langle p_{11}, p_5 | S | p_2, p_7 \rangle$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_5 + p_6 + p_{11} = p_1 + p_4 + p_8 - 2u$$

- Potential function

$$V_{\mathcal{M}}(v, w, x, y, z; m) = -\text{Li}_2\left(\frac{w}{v}\right) - \text{Li}_2\left(\frac{m^4 v}{y}\right) + \text{Li}_2\left(\frac{x}{y}\right) - \text{Li}_2\left(\frac{x}{z^2}\right) \\ + \text{Li}_2\left(\frac{w}{z}\right) - \text{Li}_2\left(\frac{z}{m^2 v}\right) + \frac{\pi^2}{3} - 2(\log(m^2))^2 \\ + \log(m^2) \log\left(\frac{y^2 z}{v^3 w}\right) + \log v \log\left(\frac{yz}{v}\right) + \log x \log\left(\frac{wz^3}{x}\right) - \log z \log(yz^2 w)$$

- Hyperbolic volume

$$\text{Im } V_{\mathcal{M}}(v, w, x, y, z; 1) = -D\left(\frac{w}{v}\right) - D\left(\frac{v}{y}\right) + D\left(\frac{x}{y}\right) - D\left(\frac{x}{z^2}\right) + D\left(\frac{w}{z}\right) - D\left(\frac{z}{v}\right)$$


with

$$\begin{pmatrix} v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1.18608 + 0.874646 i \\ 1.0 \\ -1.09737 + 0.230836 i \\ -1.23271 + 1.09381 i \\ 0.40897 - 0.337176 i \end{pmatrix}$$

- A-polynomial

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 10 & -7 & 3 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & -24 & 16 & -3 & 6 & -2 & 1 \\ 0 & 0 & 0 & 2 & -4 & -5 & 5 & -19 & 1 & 1 & 0 \\ 0 & 0 & 0 & -10 & 7 & 47 & -39 & 13 & -12 & -6 & 0 \\ 0 & 0 & -1 & 15 & 5 & 18 & -14 & 22 & -5 & 6 & 0 \\ 0 & -1 & 3 & 3 & -36 & -21 & 45 & 12 & 9 & 0 & 0 \\ 0 & 1 & 7 & -35 & 16 & -69 & 30 & 2 & -6 & 0 & 0 \\ 0 & 0 & -19 & 15 & 44 & -45 & 6 & -40 & -7 & 0 & 0 \\ 0 & 0 & 7 & 40 & -6 & 45 & -44 & -15 & 19 & 0 & 0 \\ 0 & 0 & 6 & -2 & -30 & 69 & -16 & 35 & -7 & -1 & 0 \\ 0 & 0 & -9 & -12 & -45 & 21 & 36 & -3 & -3 & 1 & 0 \\ 0 & -6 & 5 & -22 & 14 & -18 & -5 & -15 & 1 & 0 & 0 \\ 0 & 6 & 12 & -13 & 39 & -47 & -7 & 10 & 0 & 0 & 0 \\ 0 & -1 & -1 & 19 & -5 & 5 & 4 & -2 & 0 & 0 & 0 \\ -1 & 2 & -6 & 3 & -16 & 24 & -1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & -3 & 7 & -10 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

A.2.9 $K_{6_{22}}$



$$\text{Vol}(S^3 \setminus \mathcal{K}) = 4.7698896 \dots$$

- Quantum invariant

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \, \delta_C(\mathbf{p}; u) & \langle p_6, p_1 | S^{-1} | p_1, p_2 \rangle \langle p_2, p_{11} | S^{-1} | p_3, p_7 \rangle \langle p_{10}, p_7 | S | p_{12}, p_{10} \rangle \\ & \times \langle p_3, p_9 | S^{-1} | p_4, p_{11} \rangle \langle p_5, p_4 | S | p_9, p_8 \rangle \langle p_8, p_{12} | S | p_6, p_5 \rangle \end{aligned}$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_4 + p_5 + p_7 = p_8 + p_9 + p_{12} - 2u$$

- Potential function


$$\begin{aligned}
 V_{\mathcal{M}}(v, w, x, y, z; m) = & -\operatorname{Li}_2\left(\frac{w}{v^2}\right) - \operatorname{Li}_2\left(\frac{m^2 v}{w^2 x^2}\right) + \operatorname{Li}_2\left(\frac{m^2 v}{y}\right) + \operatorname{Li}_2\left(\frac{y}{v^2 x}\right) \\
 & - \operatorname{Li}_2\left(\frac{v x z}{w}\right) + \operatorname{Li}_2\left(\frac{x z}{y}\right) - \left(\log(m^2)\right)^2 + \log v \log(w z^2) - 3(\log w)^2 \\
 & + \log(m^2) \log\left(\frac{w^3 x^4 z^2}{y}\right) - 2 \log y \log(v^2 x^2 y z) + \log x \log\left(\frac{v^8}{w^4 x}\right)
 \end{aligned}$$

- Hyperbolic volume

$$\operatorname{Im} V_{\mathcal{M}}(v, w, x, y, z; 1) = -D\left(\frac{w}{v^2}\right) - D\left(\frac{v}{w^2 x^2}\right) + D\left(\frac{v}{y}\right) + D\left(\frac{y}{v^2 x}\right) - D\left(\frac{v x z}{w}\right) + D\left(\frac{x z}{y}\right)$$

with

$$\begin{pmatrix} v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0.127216 + 0.708358 i \\ -0.534645 + 1.01011 i \\ -0.40328 + 0.755585 i \\ 0.66186 - 0.301752 i \\ -0.54977 - 1.03005 i \end{pmatrix}$$

A.2.10 $K6_{33}$ or 10_{140} 

$$\text{Vol}(S^3 \setminus \mathcal{K}) = 5.212567 \dots$$

- Quantum invariant

$$\begin{aligned} Z_\gamma(\mathcal{M}_u) = \int_{\mathbb{R}} d\mathbf{p} \, \delta_C(\mathbf{p}; u) \langle p_1, p_7 | S | p_5, p_3 \rangle \langle p_2, p_9 | S^{-1} | p_1, p_4 \rangle \langle p_3, p_{12} | S^{-1} | p_{10}, p_2 \rangle \\ \times \langle p_5, p_4 | S^{-1} | p_6, p_{11} \rangle \langle p_8, p_6 | S^{-1} | p_7, p_{12} \rangle \langle p_{10}, p_{11} | S^{-1} | p_9, p_8 \rangle \end{aligned}$$

- Condition $\delta_C(\mathbf{p}; u)$

$$p_3 + p_6 + p_{11} = p_2 + p_7 + p_9 - 2u$$

- Potential function

$$\begin{aligned} V_{\mathcal{M}}(v, w, x, y, z; m) = -\text{Li}_2\left(\frac{1}{vy}\right) - \text{Li}_2\left(\frac{1}{m^2 wy}\right) + \text{Li}_2\left(\frac{w^2 x}{y}\right) \\ - \text{Li}_2(v m^2 w y) - \text{Li}_2(v m^2 w x z) - \text{Li}_2(m^2 y z) + \frac{2\pi^2}{3} \\ - (\log(v m^2))^2 - 2 \log v \log w - 2 \log(m^2) \log(w y) \\ - \log v \log(x y) - \log(m^2) \log z - \log y \log(z w y) \end{aligned}$$

- Hyperbolic volume

$$\begin{aligned} \text{Im } V_{\mathcal{M}}(v, w, x, y, z; 1) = -D\left(\frac{1}{vy}\right) - D\left(\frac{1}{wy}\right) + D\left(\frac{w^2 x}{y}\right) \\ - D(v w y) - D(v w x z) - D(y z) \end{aligned}$$

with

$$\begin{pmatrix} v \\ w \\ x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -1.1238 - 0.998279 i \\ -0.439261 - 0.570751 i \\ -0.836795 + 1.7323 i \\ -0.829546 - 0.0564355 i \\ -0.549394 + 0.740149 i \end{pmatrix}$$

• A-polynomial

$$\begin{pmatrix} 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -9 & 0 & 0 & 0 & 0 & 0 \\ 0 & -2 & 8 & -12 & -2 & 0 & 0 & 0 \\ 0 & 5 & -4 & 6 & -12 & -3 & 0 & 0 \\ 0 & -6 & 5 & -7 & 4 & -13 & -6 & -1 \\ 0 & 4 & -7 & 8 & -20 & 1 & -2 & 0 \\ 0 & -2 & 1 & -20 & 8 & -7 & 4 & 0 \\ -1 & -6 & -13 & 4 & -7 & 5 & -6 & 0 \\ 0 & 0 & -3 & -12 & 6 & -4 & 5 & 0 \\ 0 & 0 & 0 & -2 & -12 & 8 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 & -9 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & -1 & 0 \end{pmatrix}$$

R

- [1] E. W. Barnes, *Theory of the double gamma functions*, Phil. Trans. Roy. Soc. A **196**, 265–388 (1901).
- [2] V. V. Bazhanov and N. Reshetikhin, *Remarks on the quantum dilogarithm*, J. Phys. A: Math. Gen. **28**, 2217–2226 (1995).
- [3] R. Benedetti and C. Petronio, *Lectures on Hyperbolic Geometry*, Universitext, Springer-Verlag, Berlin, 1992.
- [4] S. Betley, J. H. Przytycki, and T. Żukowski, *Hyperbolic structures on Dehn filling of some punctured-torus bundles over S^1* , Kobe J. Math. **3**, 117–147 (1986).
- [5] A. G. Bytsko and J. Teschner, *R-operator, co-product and Haar-measure for the modular double of $U_q(sl(2, R))$* , Commun. Math. Phys. **240**, 171–196 (2003).
- [6] P. J. Callahan, J. C. Dean, and J. R. Weeks, *The simplest hyperbolic knots*, J. Knot Theory and Its Ramifications **8**, 279–297 (1999).
- [7] L. Chekhov and V. V. Fock, *Quantum Teichmüller space*, Theor. Math. Phys. **120**, 1245–1259 (1999).
- [8] L. O. Chekhov and R. C. Penner, *Introduction to quantum Thurston theory*, Russ. Math. Surveys **58**, 1141–1183 (2003).
- [9] D. Cooper, M. Culler, H. Gillet, D. D. Long, and P. B. Shalen, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. **118**, 47–84 (1994).
- [10] D. Cooper and D. D. Long, *Representation theory and the A-polynomial of a knot — knot theory and its applications*, Chaos, Solitons & Fractals **9**, 749–763 (1998).
- [11] D. Coulson, O. A. Goodman, C. D. Hodgson, and W. D. Neumann, *Computing arithmetic invariants of 3-manifolds*, Experimental Math. **9**, 127–152 (2000).
- [12] L. D. Faddeev, *Modular double of quantum group*, in G. Dito and D. Sternheimer, eds., *Conference Mosh Flato 1999 — Vol. I. Quantization, Deformations, and Symmetries*, vol. 21 of *Math. Phys. Stud.*, pp. 149–156, Kluwer, Dordrecht, 2000.
- [13] L. D. Faddeev and R. M. Kashaev, *Quantum dilogarithm*, Mod. Phys. Lett. A **9**, 427–434 (1994).
- [14] L. D. Faddeev, R. M. Kashaev, and A. Y. Volkov, *Strongly coupled quantum discrete Liouville theory I: algebraic approach and duality*, Commun. Math. Phys. **219**, 199–219 (2001).
- [15] W. Floyd and A. Hatcher, *Incompressible surfaces in punctured torus bundles*, Topology Appl. **13**, 263–282 (1982).
- [16] S. Francaviglia, *Algebraic and geometric solutions of hyperbolicity equations*, Topol. Appl. **145**, 91–118 (2004).
- [17] S. Garoufalidis and T. T. Q. Le, *The colored Jones function is q-holonomic*, Geom. Topol. **9**, 1253–1293 (2005).

- [18] S. Gukov, *Three-dimensional quantum gravity, Chern–Simons theory, and the A-polynomial*, Commun. Math. Phys. **255**, 577–627 (2005).
- [19] K. Hikami, *Hyperbolic structure arising from a knot invariant*, Int. J. Mod. Phys. A **16**, 3309–3333 (2001).
- [20] ———, *Hyperbolicity of partition function and quantum gravity*, Nucl. Phys. B **616**, 537–548 (2001).
- [21] ———, *Hyperbolic structure arising from a knot invariant II. completeness*, Int. J. Mod. Phys. B **16**, 1963–1970 (2002).
- [22] ———, *Asymptotics of the colored Jones polynomial and the A-polynomial: Twist knot and torus knot*, preprint (2004).
- [23] ———, *Difference equation of the colored Jones polynomial for the torus knot*, Int. J. Math. **15**, 959–965 (2004).
- [24] J. Hoste and M. B. Thistlethwaite, *Knotscape*, <http://www.math.utk.edu/~morwen/> (1999).
- [25] V. F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, Bull. Amer. Math. Soc. **12**, 103–111 (1985).
- [26] R. M. Kashaev, *Quantum dilogarithm as a 6j-symbol*, Mod. Phys. Lett. A **9**, 3757–3768 (1994).
- [27] ———, *An invariant of triangulated links from quantum dilogarithm*, POMI **224**, 208–214 (1995).
- [28] ———, *A link invariant from quantum dilogarithm*, Mod. Phys. Lett. A **10**, 1409–1418 (1995).
- [29] ———, *The hyperbolic volume of knots from quantum dilogarithm*, Lett. Math. Phys. **39**, 269–275 (1997).
- [30] ———, *On the spectrum of Dehn twists in quantum Teichmüller theory*, in A. N. Kirillov and N. Liskova, eds., *Physics and Combinatorics — Proceedings of the Nagoya 2000 International Workshop*, pp. 63–81, World Scientific, Singapore, 2001.
- [31] S. Kharchev, D. Lebedev, and M. Semenov-Tian-Shansky, *Unitary representations of $U_q(\mathfrak{sl}(2, \mathbb{R}))$, the modular double, and the multiparticle q -deformed Toda chains*, Commun. Math. Phys. **225**, 573–609 (2002).
- [32] A. N. Kirillov, *Dilogarithm identities*, Prog. Theor. Phys. Suppl. **118**, 61–142 (1995).
- [33] A. N. Kirillov and N. Y. Reshetikhin, *Representations of the algebra $U_q(\mathfrak{sl}(2))$, q -orthogonal polynomials and invariants of links*, in V. G. Kac, ed., *Infinite Dimensional Lie Algebra and Groups*, pp. 285–339, World Scientific, Singapore, 1989.
- [34] N. Kurokawa, *Multiple sine functions and Selberg zeta functions*, Proc. Japan Acad. Ser. A Math. Sci **67**, 61–64 (1991).
- [35] T. T. Q. Le, *The colored Jones polynomial and the A-polynomial of two-bridge knots*, math.GT/0407521 (2004).
- [36] L. Lewin, *Polylogarithms and Associated Functions*, North-Holland, New York, 1981.
- [37] R. Meyerhoff, *Hyperbolic 3-manifolds with equal volumes but different Chern–Simons invariants*, in D. B. A. Epstein, ed., *Low-Dimensional Topology and Kleinian Groups*, vol. 112 of *London Math. Soc. Lecture Notes*, pp. 209–215, Cambridge Univ. Press, Cambridge, 1986.
- [38] J. Milnor, *Hyperbolic geometry: the first 150 years*, Bull. Amer. Math. Soc. **6**, 9–24 (1982).
- [39] H. Murakami, *Some limits of the colored Jones polynomials of the figure-eight knot*, Kyungpook Math. J. **44**, 369–383 (2004).
- [40] ———, *A version of the volume conjecture*, math.GT/0603217 (2006).
- [41] H. Murakami and J. Murakami, *The colored Jones polynomials and the simplicial volume of a knot*, Acta Math. **186**, 85–104 (2001).
- [42] W. D. Neumann and J. Yang, *Invariants from triangulations of hyperbolic 3-manifolds*, Electron. Res. Announc. Amer. Math. Soc. **1**, 72–79 (1995).
- [43] ———, *Bloch invariants of hyperbolic 3-manifolds*, Duke Math. J. **96**, 29–59 (1999).
- [44] W. Z. Neumann and D. Zagier, *Volumes of hyperbolic three-manifolds*, Topology **24**, 307–332 (1985).
- [45] G. Ponzano and T. Regge, *Semiclassical limit of Racah coefficients*, in F. Bloch, ed., *Spectroscopic and Group Theoretical Methods in Physics*, pp. 1–58, North-Holland, 1968.
- [46] S. N. M. Ruijsenaars, *First order analytic difference equations and integrable quantum systems*, J. Math. Phys. **38**, 1069–1146 (1997).
- [47] R. Scharein, *Knotplot*, <http://www.pims.math.ca/knotplot/>.

- [48] T. Shintani, *On a Kronecker limit formula for real quadratic fields*, J. Fac. Sci. Univ. Tokyo **24**, 167–199 (1977).
- [49] M. Takahashi, *On the concrete construction of hyperbolic structures of 3-manifolds*, Tsukuba J. Math. **9**, 41–83 (1985).
- [50] W. P. Thurston, *The geometry and topology of three-manifolds*, Lecture Notes in Princeton University, Princeton (1980).
- [51] ———, *Hyperbolic structures on 3-manifolds. I. deformation of acylindrical manifolds*, Ann. Math. **124**, 203–246 (1986).
- [52] J. Weeks, *SnapPea*, <http://thames.northnet.org/weeks/>.
- [53] E. Witten, *Quantum field theory and the Jones polynomial*, Commun. Math. Phys. **121**, 351–399 (1989).
- [54] S. L. Woronowicz, *Quantum exponential function*, Rev. Math. Phys. **12**, 873–920 (2000).
- [55] T. Yoshida, *The η -invariant of hyperbolic 3-manifolds*, Invent. Math. **81**, 473–514 (1985).

E-mail address: hikami@phys.s.u-tokyo.ac.jp

URL: <http://gogh.phys.s.u-tokyo.ac.jp/~hikami/>

D P , G S S „ U T „ H 7–3–1, B , T
113–0033, J .